

Contents lists available at BioMedSciDirect Publications

International Journal of Biological & Medical Research

Journal homepage: www.biomedscidirect.com



Original Article

Effect of Fast Tempo Songs on Heart Rate of Young Adults

Manibalan Vijayaraman a and Gunasekaran Ramanathan b

Department of Physiology, Vinayaka Mission Medical College and Research Institute, Karikal , India. Department of Physiology, VMKV Medical College, Vinayaka Missions University, Salem, TN, India

ARTICLE INFO

Keywords: Fast tempo Slow tempo Song HRMusic

ABSTRACT

Earlier studies show listening music has effect on heart rate however, the rationale of the present study was to find whether the different kind of tempo songs have affect on heart rate (HR). For these study normal, healthy medical students 19-21 years, both sexes were selected (N30). In four different stages their HR, was measured 1) Baseline (without any stimulus) 2) before exercise (songs played) 2) after 5-6 min exercise using Bi-cycle Ergo meter (no song played) 4) after exercise, slow tempo 60-70 beats/ min, medium 80-90 beats/ min and fast tempo 120-160 beats / min songs were played (2.5-3 min) with a resting period of 30 min between one tempo to another tempo song. The result shows that the fast tempo songs significantly (P<0.005) increases the HR, when compare to other tempo songs it indicate that fast tempo song has effect on HR

© Copyright 2010 BioMedSciDirect Publications IJBMR - ISSN: 0976:6685. All rights reserved.

Introduction

Music is the big part of human life. It's competent of numerous health benefits including lowering stress levels, raising states of consciousness and changing moods. Numerous studies reported that music has favorable physiological effect on human especially during stress, anxiety and depression [1, 2]. Listening music that imitate or improves their emotional state [3, 4], manipulate emotions [5] decreasing HR, and blood pressure (BP), and pain reduction [6]. However; it's mainly based on the fast or slow tempo of music, extreme music leads to emotions, anger and aggression [7] tension [8] leads stress. Most of the earlier studies showed that music has positive effect but only limited study shows the negative effect of music [9] hence, this study has been undertaken. The purpose of the present study is to assess whether the different music tempo has direct effect on increasing HR in normal adults.

Materials and Methods

All methods were carried out in accordance with the approved guidelines from the VMKV medical university institutional ethical board. Normal healthy young adults, both sex, age between [19-22 years] normal BMI (N30). Each subject signed informed consent was obtained.

Song selection and analysis

An analysis of the 500 different language songs, we finalized based on participant like and dislike or anger creating songs from that slow tempo 60-70 beats/min, medium 80-90 beats/min and fast tempo 120-160 beats / min songs were selected. Each Song was analyzed beats per min (BPM) by Piston soft BPM detector (online)

Methods

The selected subjects were in four different stages their HR, was measured 1) Initially baseline (normal room temperature, without any stimulus) 2) before exercise (songs played) 3) after 5-6 min exercise using Bi-cycle Ergo meter (no song played) 4) after exercise. slow tempo 60-70 beats/min, medium 80-90 beats/min and fast tempo 120-160 beats / min songs were played (2.5-3 min) with a resting period of 30 min between one tempo to another tempo song. The songs were delivered through the head phone and sound intensity was adjusted according to the subject's tolerance. For HR measurement computerized Electrocardiogram (ECG) was used and R-R interval calculated.

HR Measurement

Computerized ECG recorder (CARDIOART 8408 VIEW, India) was used. Based on experimental procedure each group subjects were allowed to record continuous ECG for 4-5 min automatically, it shows the mean HR/min

Moderate exercise

After baseline measurements subjects were asked to do moderate exercise using Bi-cycle Ergo meter (INCO, India) for 5 min (distance covered / unit time is constant) after that very small break 2-3 min rest and the subjects HR was measured

Statistical analysis

Descriptive data are expressed as mean ± standard deviation (SD). One-way ANOVA was used to analyze the data. The level of significance was P < 0.05. All data were analyzed using SPSS for Windows version 17.0 (SPSS Inc., Chicago, IL, USA).

Result

We evaluated heart function from HR. The data were collected from 30 healthy subjects. The effect of different tempo songs on

^{*} Corresponding Author: Dr. Gunasekaran Ramanathan PhD, PDF (USA) Professor, Department of Physiology VMKV Mediacal College, Seeragapadi, Salem, India Email: gunasekaranrr@gmail.com

heart function was analyzed in normal condition (before exercise) (Table 1.) show the mean and standard deviation (SD) of HR. For comparison study one way ANOVA was used, our result shows that fast tempo songs has significantly (P<0.01) increase the HR. when compare to baseline (no song played) and slow or medium tempo songs (Table 2.).

Table 1.Effect of different tempo songs on HR in normal condition (Before exercise)

	Baseline Mean ±SD	Before exercise (Song played) Mean ±SD		
	no song plaved	Slow tempo	Medium tempo	Fast tempo
Heart rate (HR)	71.60 ±.4.174	67.60±2.313	73.40 ±7.093	80.87±4.754

Note: Mean of normal healthy subjects (N30)

Table 2.Comparition of deferent tempo songs on HR in normal condition (Before exercise)

	Heart rate HR		
Multiple comparison	Mean difference	P value	
Base line vs Fast tempo song	-9.267	P < 0.001	
Slow tempo vs Fast tempo song	-13.27	P < 0.001	
Base line vs Medium tempo song	-1.800	P > 0.05	

Note: P value significant P < 0.001, (N30)

Table 3. Effect of different tempo songs on HR after exercise

	Baseline Mean ±.SD	After exercise (no song played) Mean ±SD	After exercise (song played) Mean ±SD		
	no song played	no song played	Slow tempo	Medium tempo	Fast tempo
Heart rate (HR)	71.60 ±.4.174	103.1±12.78	97.93±12.8 8	105.4±12.4 1	113.4±9.77 5

Note: Mean of normal healthy subjects (N30)

Table 4. Comparison of deferent tempo songs on HR after exercise

	Heart rate HR		
Multiple comparison	Mean difference	P value	
After exercise (no song) vs Fast tempo song	-10.33	P < 0.01	
After exercise Slow tempo vs Fast tempo song	-15.47	P < 0.001	
After exercise (no song) vs Medium tempo song	-2.300	P > 0.05	

Note: P value significant P < 0.05 and P < 0.001 (N30)

Further, to confirm the fast tempo songs effect on heart function, we analyzed the effect of different tempo songs on after exercise, our result shows that even after exercise the fast tempo songs has significantly (P<0.05) increases the HR, when compare to base line or after exercise (no song played) or other tempo songs played (Table 3 and 4)

Discussion

The aim of the study is to find the negative effect music hearing, our study result shows that fast tempo songs before and after exercise significantly increases the HR. It indicates fast tempo song has negative effect on heart function, our result consistent with the previous study report rap music, linked with fast tempo music, causes increasing HR [10], extreme music causes anger and promotes aggressive behavior (11, 12, 13). Recent studies also shows that extreme music leads to anger, such as aggression [9] and stress it could increases the sympathetic tone due to activation of HPA [14 15] which may increases the HR, [16, 17]. Earlier studies hypothesized that there would be a significant effect of different tempo music on sympathetic nervous system level of activation [18]. Our findings support the earlier concept

Conclusion

The current study conclude that fast tempo music increase the activity of HR it may due to anger, aggression and stress via increase sympathetic tone. However, further research is required to replicate these findings and long term effects of fast tempo songs.

Acknowledgments

This study was financially supported by Vinayaka Missions University (VMU). Hereby, the author would like thanks to the young medical students who were involved in this study and to Dr .H.S. Gulapalli, HOD, Department of physiology for kind support and Mrs. Bama for technical support Vinayaka Mission Medical College and Research Insatiate, Karikal, India

REFERENCES

- Bernardi L, Porta C, Casucci G, Balsamo R, Bernardi NF, Fogari R, et al. Dynamic interactions between musical, cardiovascular, and cerebral rhythms in humans. Circulation. 2009; 119 (25):3171-80. PMID: 19569263
- Nyklicek I, Thayer JF, Van Doornen LJP. Cardio respiratory differentiation of musically induced emotions. J Psychophysiol. 1997; 11:304-321.
- Saarikallio S. Music as emotional self-regulation throughout adult hood. Psychol Music. 2011; 39 (no 3): 307-327. doi: 10.1177/0305735610374894
- Thoma MV, Ryf S, Mohiyeddini C, Ehlert U, Nater UM. Emotion regulation through listening to music in everyday situations. Cogn Emot. 2012; 26(3): 550-60. PMID: 19834599
- Juslin PN, Sloboda J. Hand book of music and emotion: Theory, research, applications. Oxford: Oxford University Press; 2010.
- Roy M, Peretz I, Rainville P. Emotional valence contributes to music-induced analgesia. Pain 2008; 134, 140–147. PMID: 17532141
- Labbé E, Schmidt N, Babin J, Pharr M. Coping with stress: the effectiveness of different types of music. Appl Psychophysiol Bio feedback. 2007; 32: 163–168. PMID: 17965934
- Krumhansl, Carol L. Music: A link between cognition and emotion. Current Directions of Psychological Science. 2002; 11(2): 45-40.

- Selfhout MH, Delsing M.J, terBogt TF, Meeus WH. Heavy metal and hiphopstyle preferences and externalizing problem behavior-at wo-wave longitudinal study. Youth Soc.2008 39 (no 4): 435-452. doi: 10.1177/0044118X07308069
- 10. Fairclough SH, van der Zwaag M, Spiridon E, Westerink J. Effects of mood induction via music on cardiovascular measures of negative emotion during simulated driving. Physiol Behav. 2014; 22(129):173-80. PMID: 24603216
- 11. Sharman L, Dingle GA. Extreme metal music and anger processing. Front Hum Neurosci. 2015; 21(9): 272. PMID: 26052277
- 12. Arnett,j. Metal heads: alienation heavy metal music and adolescent. Boulder: West view Press; 1996.
- 13. Gowensmith WN, Bloom LJ. The effect soft heavy metal music on arousal and anger. J.MusicTher.1997; 34(1): 33-45. doi: 10.1093/jmt/34.1.33
- York: Springer Verlag Publishing Company.
 Tsigos C, Chrousos GP Hypothalamic-pituitary-adrenal axis,neuroendocrine factors and stress. J Psychosom Res 2002; 53: 865-871. PubMed: 12377295.
- Sapolsky RM, Romero M, Munck AU How do Glucocorticoids influence Stress Responses? Integrating Permissive, Suppressive, Stimulatory, and Preparative Actions? Endocr Rev 2000. 21: 55-89. PubMed: 10696570.

- 16. Neus Herrero, Marien Gadea, Gabriel Rodríguez-Alarcón, Raúl Espert, Alicia Salvador. What happens when we get angry? Hormonal, cardiovascular and asymmetrical brain responses. Hormones and Behavior.2010;57(3):276-83.
- Betensky JD, Contrada RJ. Depressive symptoms, trait aggression, and cardio vascular reactivity to a laboratory stressor. Ann Behav Med. 2010; 39(2):184-91. PMID: 20379807
- Sutoo D, Akiyama K. Music improves dopaminergic neurotransmission: demonstration based on the effect of music on blood pressure regulation. Brain Res. 2004; 1016 (2):255-62. PMID: 15246862

© Copyright 2010 BioMedSciDirect Publications IJBMR - ISSN: 0976:6685.
All rights reserved.