Int J Biol Med Res. 2025; 16(1): 7930-7932

Contents lists available at BioMedSciDirect Publications

International Journal of Biological & Medical Research

Journal homepage: www.biomedscidirect.com

Original Article

Exploring Fermentation Rate Variability: A Comparative Study of Fresh and Commercial Fruit and Vegetable Juices

Dr. S. P. Patil*, Miss S. S.Nejkar, Miss F. A.Mutwalli, Mr. C. A. Mujawar, Miss R. R. Lapate

Department of Pharmaceutical Chemistry, Dr. Shivajirao Kadam College of Pharmacy,

Kasabe Digraj, Sangli, Maharashtra, 416 305, India.

ARTICLEINFO

Keywords:
Rates ofFermentation,
fruit juice,
vegetable juice,
baker'syeast,
Pasteur's salt,
Fehling's test.

ABSTRACT

This research focused on examining the fermentation rates of a variety of fruits, vegetables, and commercially available fruit juices. The fruits selected for this study included apple, watermelon, orange, mango, pomegranate, and grapes, while the vegetables used were carrot, bitter gourd, bottle gourd, spinach, and beetroot. Each of these juices was subjected to fermentation under identical conditions with the help of yeast. To monitor the progress of fermentation, Fehling's solution was used to test the presence of sugars. The time taken for the complete fermentation process was recorded by noting when the solution no longer gave a negative result, indicating that fermentation was finished. The primary objective of this study was to compare and analyse the fermentation rates across the different fruit and vegetable juices.

© Copyright 2023 BioMedSciDirect Publications IJBMR -ISSN: 0976:6685.

Introduction

This study presents a comparative analysis of the fermentation process between freshly extracted fruit and vegetable juices, as well as commercially available fruit juices. The term "fermentation" is derived from the Latin word "fermentare," meaning "to leaven" or "to sour [1]."The study primarily focuses on the varying sugar content present in both extracted and commercial juices, examining how these differences impact the time required for fermentation. Understanding the duration and process of fermentation is crucial, as fermentation has long been employed for multiple purposes, including food preservation, product development (e.g., cheese, yogurt, beer), and enhancing flavor. Fermentation is an ancient practice deeply integrated into daily life, seen in foods like dosa and idli batter in Indian cuisine [2].

Fermentation, at its core, is a metabolic process wherein microorganisms such as yeasts or bacteria break down sugars into simpler compounds, such as alcohol, organic acids, and gases. The process begins when these microorganisms come into contact with sugars or carbohydrates in a suitable environment. Through enzymatic reactions, complex sugars are decomposed into simpler molecules. For instance, in alcoholic fermentation, yeast converts glucose into ethanol and carbon dioxide. The by-products of fermentation—alcohol, acids, and gases—vary depending on the type of microorganism involved[3-4].

Corresponding author:

Dr. Sarika Pranam Patil
Department of Pharmaceutical Chemistry,
Dr. Shivajirao Kadam College of Pharmacy, Kasabe Digraj, Sangli, Maharashtra, 416 305, India.
Email: naradesarika@gmail.com

© Copyright 2023 BioMedSciDirect Publications IJBMR -ISSN: 0976:6685.

As an anaerobic process, fermentation allows microorganisms to generate energy through pathways that do not require oxygen, enabling their survival in oxygen-deficient environments. The outcome of fermentation is influenced by several factors, including the type of microorganism used, the starting material, temperature, pH levels, nutrient availability, and oxygen levels. These variables not only affect the efficiency and speed of fermentation but also the quality and characteristics of the final product[5-7].

Fermentation of Fruit and Vegetable Juices

Fruit juices are naturally sweet due to their fructose and sucrose content, with fructose being the sweetest naturally occurring carbohydrate, approximately 1.2 to 1.8 times sweeter than sucrose. The fermentation of fruit juice involves the conversion of these sugars into alcohol and carbon dioxide through the action of yeast [8-10]. The process begins by selecting ripe, high-quality fruits without spoilage, washing them to remove contaminants, and extracting their juice through juicing or pressing. The extracted juice is then strained and combined with yeast to initiate fermentation. This mixture is placed in a sealed vessel equipped with an airlock to prevent oxygen exposure, allowing the yeast to metabolize the sugars. As the yeast ferments the sugars, alcohol and carbon dioxide are produced, resulting in a transformation of the juice. The same fermentation process can be applied to vegetable juices, although fruits generally contain higher sugar levels compared to vegetables. Both fruit and vegetable juices undergo similar preparation: juice extraction, yeast addition, and fermentation in a sealed container, with regular monitoring [11-13].

Chemical Composition of Fruits and Vegetables

Fruits are rich in water-soluble compounds such as sugars, organic acids, pectin, and tannins, while insoluble substances like cellulose, hemicellulose, and starch are also present. Additionally, fruits of aromatic plants contain essential vitamins like ascorbic acid (Vitamin C) and tocopherols (Vitamin E), along with reducing sugars like glucose

and fructose. They are also a source of polyunsaturated fatty acids such as linoleic and α-linolenic acids.

Vegetables, on the other hand, are nutrient-dense and low in calories, with most species comprising 75-95% water content. They provide essential biomolecules such as lipids, proteins, and carbohydrates, along with a wide range of vitamins, minerals, and phytonutrients. Vegetables contain carbohydrates like sucrose, starch, cellulose, and pectin, but their sucrose content diminishes as they ripen, being converted into glucose and fructose, which are less sweet than sucrose. This distinct chemical composition of fruits and vegetables influences both their nutritional value and the dynamics of their fermentation processes [14-17].

This study aimed to investigate the rate of fermentation across various fruits, vegetables, and commercially available fruit juices. The selected fruits included apple, watermelon, orange, mango, pomegranate, and grapes, while the vegetables chosen were carrot, bitter gourd, bottle gourd, spinach, and beetroot. Each juice was fermented under controlled conditions using yeast, and the progress of fermentation was measured using Fehling's solution. The time taken for complete fermentation was recorded for each juice, with the goal of comparing the fermentation rates across different samples. This research offers insights into how the composition of different fruits and vegetables influences the fermentation process.

MATERIALS AND METHODS

Materials:

Fresh fruits, vegetables and commercial juices were purchased from local market at Sangli city, Maharashtra, India. The research experiment was carried out using deionized double distilled water. All other chemicals and solvents were of analytical grade.

Methods:

A clean 250 ml conical flask was used to contain 5.0 ml of freshly prepared apple juice, which was then diluted with 50 ml of distilled water. To this solution, 2.0 grams of Baker's yeast and 5.0 ml of Pasteur's salt solution were added. The mixture was thoroughly shaken to ensure proper mixing of all components. The temperature of the reaction mixture was carefully maintained between 35°C and 40°C throughout the process.

After 10 minutes, 5 drops of the reaction mixture were carefully extracted from the flask and introduced into a test tube containing 2 ml of Fehling's reagent. The test tube was then placed in a boiling water bath for approximately 2 minutes. The color change or formation of precipitate was observed and recorded. This procedure was repeated at 10-minute intervals until the reaction mixture ceased to produce any red color or precipitate, indicating the completion of fermentation. The total time taken for the fermentation process to complete was noted[18].

Preparation of Pasteur's Salt Solution:

The Pasteur salt solution was prepared by dissolving 10.0 g of ammonium tartrate, 2.0 g of potassium phosphate, 0.2 g of calcium phosphate, and 0.2 g of magnesium sulphate in 860 ml of distilled water, ensuring all components were fully dissolved before use [19].

RESULTS AND DISCUSSION:

The fermentation times for various fruit and vegetable juices were measured and recorded, highlighting significant variations based on the type of juice. Among the fruit juices, apple, orange, and pomegranate required the same time for complete fermentation, each taking approximately 1.5 hours. Watermelon and grape juices took slightly longer, requiring 2 hours to complete the fermentation process. Mango juice demonstrated the longest fermentation time at 2.5 hours, possibly due to its higher sugar content and thicker consistency, which could slow down the yeast's metabolic activity.

For vegetable juices, the fermentation process was noticeably quicker compared to fruits, with bitter gourd and spinach showing the fastest completion time of just 10 minutes, likely due to their lower sugar content and simpler chemical composition. Carrot juice took 30 minutes, while cucumber juice required 40 minutes to fully ferment. The longest times were observed in bottle gourd and beetroot juices, taking 50 minutes and 1 hour, respectively. The shorter fermentation times for vegetable juices compared to fruit juices can be attributed to their lower sugar concentrations and the presence of other compounds that may influence the fermentation kinetics.

These results highlight that sugar content, juice composition, and possibly pH levels significantly affect the rate of fermentation in different fruits and vegetables. The faster fermentation times for vegetables suggest that low-sugar environments may allow yeast to metabolize the available sugars more quickly, while fruit juices with higher sugar concentrations require longer fermentation times [20-21].

In contrast to fresh fruit and vegetable juices, the fermentation times for marketed fruit juices were significantly longer. This could be attributed to the presence of preservatives, added sugars, and other stabilizers, which likely interfere with or slow down the fermentation process [22]. The time required for complete fermentation of these commercially available juices ranged from 23.5 hours to 25 hours 40 minutes.

Among the marketed juices tested, orange juice exhibited the shortest fermentation time at 23 hours 30 minutes, while pineapple juice took the longest at 25 hours 40 minutes. Watermelon and Grapes juice took 24 hours 15 minutes and 24 hours 50 minutes, respectively, to complete fermentation, which was still substantially longer than their fresh counterparts. Mango juice required 25 hours, which aligns with the higher sugar content and more processed nature of this particular juice. These prolonged fermentation times underscore the impact of processing, preservatives, and added sugars commonly found in marketed juices, all of which might delay the activity of yeast in converting sugars into alcohol and carbon dioxide [20].

The slower fermentation in marketed juices could be due to the inhibitory effects of artificial ingredients or the increased complexity of sugar structures present in processed products. This indicates that fresh juices, with their natural sugars and simpler compositions, are more conducive to rapid fermentation compared to commercially prepared juices. Time required for complete fermentation of natural fruit juices, vegetable juices and commercial juices are shown in table no. 1, 2 and 3 respectively.

Table 1: Time required for complete fermentation of different natural fruit juices

Sr. no	Fruit juice	Time required for complete fermentation
1.	Apple	1.5 hr
2.	Watermelon	2 hr
3.	Orange	1.5 hr
4.	Mango	2.5 hr
5.	Pomegranate	1.5 hr
6.	Grapes	2 hr

Table 2: Time required for complete fermentation of different natural vegetable juices

Sr. no	Vegetable juices	Time required for complete fermentation
1.	Carrot	30 min
2.	Bitter gourd	10 min
3.	Bottle gourd	50 min
4.	Spinach	10min
5.	Beetroot	1 hr
6.	Cucumber	40 min

Table 3: Time required for complete fermentation of different commercial juices

Sr. no	Marketed juices	Time required for complete fermentation
1.	Watermelon	24 hr 15 min
2.	Orange	23 hr 30 min
3.	Mango	25 hr
4.	Grapes	24 hr 50 min
5.	Pineapple	25 hr 40 min

CONCLUSION

This study reveals significant differences in fermentation rates between fresh fruit, vegetable, and marketed juices. Fresh fruit juices exhibited moderate fermentation times, influenced by their natural sugar content, while vegetable juices fermented much faster due to lower sugar concentrations. In contrast, marketed juices demonstrated significantly longer fermentation times, likely might due to the presence of preservatives, added sugars, and other artificial components that slow yeast activity. Overall, the study highlights the impact of juice composition and processing on fermentation rates, with fresh juices being more favourable for quicker fermentation compared to their commercial counterparts.

ACKNOWLEDGEMENTS

Authors are thankful to Principal, Dr. Shivajirao Kadam College of Pharmacy,Kasabe Digraj,Sangli, Maharashtra, India for providing laboratory facilities and constant encouragement.

CONFLICT OF INTEREST

The author(s) declare(s) that they have no declaration of interests to disclose.

REFERENCES

[1] Harper, D. (n.d.). Etymology of sour. Online Etymology Dictionary. https://www.etymonline.com/word/sour

[2] Siddiqui SA, Erol Z, Rugji J, et al. An overview of fermentation in the food industry - looking back from a new perspective. Bioresour Bioprocess. 2023;10:85. doi:10.1186/s40643-023-00702-y.

[3] Maicas S. The Role of Yeasts in Fermentation Processes. Microorganisms. 2020;8(8):1142. doi:10.3390/microorganisms8081142

[4] Sharma R, Garg P, Kumar P, Bhatia SK, Kulshrestha S. Microbial Fermentation and Its Role in Quality Improvement of Fermented Foods. Fermentation. 2020; 6(4):106. https://doi.org/10.3390/fermentation6040106

[5]Buckel W. Energy Conservation in Fermentations of Anaerobic Bacteria. Front Microbiol. 2021;12:703525. doi:10.3389/fmicb.2021.703525

[6] Mengesha Y, Tebeje A, Tilahun B. A Review on Factors Influencing the Fermentation Process of Teff (Eragrostis teff) and Other Cereal-Based Ethiopian Injera. Int J Food Sci. 2022;2022:4419955. Published 2022 Mar 24. doi:10.1155/2022/4419955

[7] Sharma R, Garg P, Kumar P, Bhatia SK, Kulshrestha S. Microbial Fermentation and Its Role in Quality Improvement of Fermented Foods. Fermentation. 2020; 6(4):106. https://doi.org/10.3390/fermentation6040106

[8] Pepin A, Stanhope KL, Imbeault P. Are Fruit Juices Healthier Than Sugar-Sweetened Beverages? A Review. Nutrients. 2019;11(5):1006. Published 2019 May 2. doi:10.3390/nu11051006

[9] Monteiro-Alfredo T, Caramelo B, Arbeláez D, et al. Distinct Impact of Natural Sugars from Fruit Juices and Added Sugars on Caloric Intake, Body Weight, Glycaemia, Oxidative Stress and Glycation in Diabetic Rats. Nutrients. 2021;13(9):2956. Published 2021 Aug 25. doi:10.3390/nu13092956

[10]Pinto T, Vilela A, Cosme F. Chemical and Sensory Characteristics of Fruit Juice and Fruit Fermented Beverages and Their Consumer Acceptance. Beverages. 2022; 8(2):33. https://doi.org/10.3390/beverages8020033

[11]Galaverna G, Dall'Asta C. Production processes of orange juice and effects on antioxidant components. In: Processing and impact on antioxidants in beverages. 2014. p. 203-14. doi:10.1016/B978-0-12-404738-9.00021-0.

[12]Effect of mechanical juice extraction method on the quality of fresh-squeezed apple juice. IOP Conf Ser Mater Sci Eng. 2020;711(1):012051. doi:10.1088/1757-899X/711/1/012051.

[13]Sajjad N, Rasool A, Fazili AB, Bhat EA. Fermentation of fruits and vegetables. Plant Arch. 2020;20(Suppl 2):1338-42. e-ISSN: 2581-6063. ISSN: 0972-5210.

[14]Butnariu, M., Butu, A. (2015). Chemical Composition of Vegetables and Their Products. In: Cheung, P., Mehta, B. (eds) Handbook of Food Chemistry. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-36605-5_17

[15]https://fruitprocessingmachine.com/chemical-components-and-properties-of-fruits-and-vegetables/

[16]Molnár PJ. Fruits and vegetables. In: Food quality and standards – Vol. II. Encyclopedia of Life Support Systems (EOLSS).

[17]Ladaniya MS. Fruit quality control, evaluation, and analysis. In: Citrus fruit. 2008.

[18]https://www.scribd.com/doc/118086654/To-Study-the-rate-of-fermentation-of-fruits-and-vegetables-juices

[19]https://www.coursehero.com/file/p1v7up9a/Pasteurs-Salt-solution-Pasteur-salt-solution-is-prepared-by-dissolving-ammonium/

[20]Pepin C, Marzzacco C. The fermentation of sugars using yeast: a discovery experiment. Melbourne, FL; 2015 Apr.

[21]Thuy CX, Pham VT, Nguyen TTN, Nguyen TTN, Ton NTA, Tuu TT, Vu ND. Effect of fermentation conditions (dilution ratio, medium pH, total soluble solids, and *Saccharomyces cerevisiae* yeast ratio) on the ability to ferment cider from tamarillo (*Solanum betaceum*) fruit. Hindawi J Food Process Preserv. 2024;2024:Article ID 8841207, 17 pages. doi:10.1155/2024/8841207.

[22]Maia MS, Domingos MM, de São José JFB. Viability of Probiotic Microorganisms and the Effect of Their Addition to Fruit and Vegetable Juices. Microorganisms. 2023;11(5):1335. Published 2023 May 19. doi:10.3390/microorganisms11051335

© Copyright 2023 BioMedSciDirect Publications IJBMR -ISSN: 0976:6685.