Int J Biol Med Res. 2025; 16(3): 8180-8184

Contents lists available at BioMedSciDirect Publications

International Journal of Biological & Medical Research

Journal homepage: www.biomedscidirect.com

**Review Article**

# Immunological prospects and challenges of In vitro fertilization

**Solomon Matthias Gamde[1], Song Elizabeth Iliya[1], Odu Stephen Iyeje[2], Mohamed Umar[3], Amina Egbujo[3], Alheri Lailai[4]**

[1]Department of Medical Laboratory Science, Bingham University Karu, Nigeria
[2]Department of Internal Medicine, Federal Medical Centre, Azare, Bauchi, Nigeria
3Department of Medical Laboratory Science, University of Jos,Nigeria
[4]Department of Public Health and community Medicine, Bingham University Karu, Nigeria

ARTICLE INFO

*Keywords:*
*In Vitro Fertilization,*
*Immunological challenges,*
*Treatment of Infertility,*
*Inflammation*

ABSTRACT

In Vitro Fertilization (IVF) is the most common form of assisted reproductive technology (ART), which involves the manipulation of oocytes outside the body. Despite advances in vitro fertilization (IVF) technologies, several challenges remain, particularly those related to the immune system.The present review study will explore key immunological challenges, including immune rejection and inflammation, which affect IVF outcomes.The immune system plays a critical role in embryo implantation, placental development, and the maintenance of pregnancy. Improper regulation of immune responses can lead to implantation failure, pregnancy loss, or complications during IVF. Women who cannot use their Oocytes due to primary ovarian insufficiency (POI) or age-related decline in oocyte number can now become successfully pregnant utilizing donor oocyte IVF.While there has been significant progress in understanding and treating infertility disorders, there is still much to learn. Healthcare providers are constantly looking for new strategies and interventions to improve IVF success rates for people who have immune-related fertility issues.

© Copyright 2025 BioMedSciDirect Publications IJBMR -ISSN: 0976:6685.

## Introduction

In vitro fertilization (IVF) is one of the most widely used assisted reproductive technologies (ART) to treat infertility. Since the birth of the first IVF baby in 1978, this technique has evolved significantly and is now a standard treatment for individuals and couples facing difficulties in conceiving naturally. IVF involves the external fertilization of an egg by sperm in a laboratory setting, followed by the transfer of the resulting embryo into the uterus. Despite its widespread use, IVF success rates remain variable, with approximately 30%–40% of procedures resulting in a live birth, depending on maternal age and other factors [1].

A major factor influencing the success of IVF is the interaction between the embryo and the maternal immune system. The maternal immune system plays a pivotal role in both recognizing and tolerating the embryo, which is genetically distinct due to the presence of paternal antigens. While immune responses are crucial for defending against infections, they must be regulated during pregnancy to prevent rejection of the semi-allogeneic embryo. An imbalance in immune regulation can result in implantation failure, miscarriage, or other complications [2]. Most of the children born from IVF appear healthy, however, there is a noticeable increase in health problems, such as low birth weight, premature birth, and congenital birth defects.

Corresponding author:

**Dr. Solomon Matthias Gamde**
Department of Medical Laboratory Science,
Bingham University Karu, Nigeria
**Email id:** solomonmatthias85@gmail.com
ORCID; http://orcid/0000-0002-7631-8782

© Copyright 2025 BioMedSciDirect Publications IJBMR -ISSN: 0976:6685.

The immune system iprotectsthe body against harmful pathogens and foreign substances. It plays a critical role in maintaining the overall health and well-being of an individual. The adaptive (specific) immune system produces antibodies that are used to fight specific germs with which the body has previously come into contact. A strong, balanced immune system is vital in pregnancy as it helps the uterus accept and nurture the embryo.Embryo implantation is a highly complex process that involves a coordinated interaction between maternal immune cells and the developing embryo. The immune system helps facilitate implantation by promoting tissue remodeling and supporting a hospitable uterine environment [3].

### 2.0 MATERIALS AND METHOD

In this review published papers in an electronic database comprising Research Gate, Google Scholar, and PubMed were retrieved using search themes such as “Vitro fertilization”, “Autoimmune disorders”, “Immune modulators”, “Immunological Factors” OR “Maternal Immune System” were examined.

### 2.1 Searching strategies and sources

Articles that met the criteria were identified using keywords and MeSH words in an electronic database, and hand searching using bibliographic or reference information from identified studies.

### 2.2 Inclusion and exclusion criteria

The purpose for exclusion at the full-text screen level was documented by the PRISMA framework. Two authors independently screened articles using the inclusion criteria. Duplicate studies from electronic databases obtained through the search strategy were removed using the EndNote program. The titles and abstracts of the articles found through the search strategy were independently screened by two reviewers to eliminate obvious non-relevant papers. The full-text versions of the remaining potentially eligible articles were then retrieved and independently accessed to determine if they met the inclusion criteria.

## 3.0 RESULT AND DISCUSSION

### 3.1 Some immunological cells in IVF

Crucial immune players in IVF include the regulatory T cells (Tregs), natural killer (NK) cells, dendritic cells, and various cytokines that mediate immune tolerance [4].

1. Regulatory T Cells (Tregs) are critical in establishing immune tolerance during pregnancy, helping prevent the maternal immune system from attacking the embryo, which contains paternal antigens. Tregs suppress the activity of other immune cells which could lead to embryo rejection. Studies have demonstrated that increased Treg populations are associated with successful pregnancies, both in natural conception and IVF [5]. plays a central role in maintaining tolerance to the embryo by suppressing potential immune attacks against the paternal antigens expressed by the embryo. Tregs have been shown to increase in numbers during early pregnancy and are associated with successful implantation [5].

2. Natural Killer (NK) Cells are essential for proper placental development and tissue remodeling, their over-activation can lead to inflammation, which may hinder embryo implantation. Elevated NK cell activity has been linked to recurrent implantation failure [6].

3. Cytokines are signaling proteins that regulate immune responses, and play an essential role in the dialogue between the embryo and maternal immune system. A balance between pro-inflammatory (Th1) and anti-inflammatory (Th2) cytokines is necessary for successful implantation. An overproduction of Th1 cytokines, such as tumor necrosis factor-alpha (TNF-α), may create an inflammatory environment that is hostile to the embryo [4].

### 3.2 Immunological Challenges in IVF

Despite the immune system's essential role in promoting pregnancy, several immunological factors can present challenges to IVF success. These include recurrent implantation failure (RIF), immune dysregulation, and autoimmune conditions. For instance, women with recurrent miscarriages or RIF often exhibit abnormalities in immune function, such as excessive NK cell activity or altered cytokine profiles [7]. Autoimmune disorders, such as antiphospholipid syndrome (APS), can also interfere with embryo implantation and increase the risk of pregnancy loss [8].

Given the complex interplay between the immune system and reproductive processes, immunomodulatory therapies, such as intravenous immunoglobulin (IVIG), corticosteroids, and intralipid infusions, have been explored as potential treatments for women experiencing immune-related reproductive failures. However, these therapies require careful consideration to avoid excessive immunosuppression, which can increase infection risks and negatively impact pregnancy outcomes [6].The role of the immune system in IVF is critical to both embryo implantation and pregnancy maintenance. While immune tolerance mechanisms enable successful pregnancies, dysregulation of immune responses can lead to reproductive failures, including implantation failure and miscarriage. As research progresses, a better understanding of the immunological factors at play may lead to improved therapies and higher success rates in IVF.

#### 1. Immune Rejection of the Embryo

One of the primary immunological challenges in IVF is the risk of immune rejection of the embryo. The embryo is a semi-allograft, meaning it carries paternal antigens that are foreign to the maternal immune system. The maternal immune system must, therefore, develop mechanisms to tolerate these foreign antigens while still providing immune protection against potential pathogens.

**Failure of Immune Tolerance:** During a successful pregnancy, the maternal immune system adapts to tolerate the semi-allogeneic embryo through the action of regulatory T cells (Tregs) and the expression of non-classical HLA molecules, such as HLA-G, by the trophoblast. However, in some women undergoing IVF, this immune tolerance may not be properly established, leading to the immune system attacking and rejecting the embryo, contributing to implantation failure [5].

**Elevated NK Cell Activity:** Natural killer (NK) cells play a dual role in pregnancy: they are essential for trophoblast invasion and vascular remodeling but can also become overactive and cytotoxic, leading to embryo rejection. Elevated NK cell activity has been associated with recurrent implantation failure (RIF) and miscarriage. In IVF patients, elevated levels of cytotoxic NK cells may interfere with the establishment of immune tolerance, leading to the rejection of the transferred embryo [6].

#### 2. Inflammation and Cytokine Imbalance

Inflammation, mediated by an imbalance of cytokines, is another significant challenge in IVF. Successful implantation requires a balance between pro-inflammatory and anti-inflammatory cytokines. Dysregulation of this balance can lead to a hostile uterine environment, preventing embryo implantation.

**Pro-Inflammatory Cytokines (Th1 Response):** Cytokines such as tumor necrosis factor-alpha (TNF-α), interleukin-6 (IL-6), and interferon-gamma (IFN-γ) are part of the pro-inflammatory Th1 response. While a mild inflammatory environment is necessary for embryo implantation, excessive production of these cytokines can lead to chronic inflammation, creating a hostile uterine environment. Elevated levels of TNF-α and IFN-γ have been associated with implantation failure and recurrent miscarriage in IVF patients [4].

**Anti-Inflammatory Cytokines (Th2 Response):** Anti-inflammatory cytokines, such as interleukin-10 (IL-10) and interleukin-4 (IL-4), are part of the Th2 response, which promotes immune tolerance and a favorable environment for implantation. A deficiency in Th2 cytokine activity or a failure to shift from a pro-inflammatory (Th1) to an anti-inflammatory (Th2) environment can hinder successful implantation [5].

**Imbalance in Th1/Th2 Cytokine Ratios:** The delicate balance between Th1 and Th2 cytokines is crucial for successful implantation and pregnancy. An abnormal shift towards a Th1-dominant response has been implicated in pregnancy loss and poor IVF outcomes. Conversely, a Th2-dominant environment is more conducive to embryo implantation and the continuation of pregnancy [4].

#### 3. Autoimmune Disorders and Reproductive Failure

Autoimmune disorders are another challenge for women undergoing IVF. Autoimmune conditions can lead to the production of antibodies that attack the body's tissues, including those of the reproductive system. These antibodies can interfere with embryo implantation and increase the risk of miscarriage.

**Antiphospholipid Syndrome (APS):** APS is an autoimmune disorder characterized by the presence of antiphospholipid antibodies, which can lead to blood clot formation and placental insufficiency. Women with APS have a higher risk of recurrent miscarriage and implantation failure, and they may require anticoagulation therapy to improve IVF outcomes [8].

**Antinuclear Antibodies (ANA):** ANA are autoantibodies that target the body's cells and tissues. Elevated ANA levels have been associated with implantation failure and recurrent miscarriage, particularly in women undergoing IVF. These antibodies may interfere with the maternal immune system's ability to tolerate the embryo [7].

#### 4. Endometrial Receptivity and Immune Dysregulation

For successful embryo implantation, the endometrium must be receptive at the time of embryo transfer. Immune dysregulation in the uterine environment can disrupt this receptivity, preventing the embryo from implanting properly.

**Endometrial NK Cells:** uNK cells play a role in the remodeling of uterine blood vessels during early pregnancy. However, increased numbers of cytotoxic NK cells in the endometrium have been associated with implantation failure. In IVF, women with elevated uNK cell activity may experience repeated implantation failure due to an overly aggressive immune response in the endometrium [6].

**Chronic Endometritis:** Chronic endometritis, characterized by persistent inflammation of the endometrium, can negatively affect endometrial receptivity. Chronic inflammation, often caused by bacterial infections or immune dysregulation, has been linked to poor IVF outcomes, including implantation failure [5].

**5. Immunosuppressive Therapies and Their Risks**

While immunosuppressive therapies, such as corticosteroids, intravenous immunoglobulin (IVIG), and intralipid infusions, are often used to modulate immune responses in IVF, they come with potential risks.

**Excessive Immunosuppression:** Overuse of immunosuppressive therapies can compromise the maternal immune system's ability to defend against infections. This can lead to an increased risk of infections during pregnancy and may negatively impact both maternal and fetal health [9].

**Tailored Treatment:** Immunosuppressive therapies must be carefully tailored to each patient's unique immunological profile. Over-suppression of immune responses may lead to complications, while inadequate suppression may result in persistent immune rejection of the embryo [7].

**3.3Challenges in IVF**

Despite advances in in vitro fertilization (IVF) technologies, several challenges remain, particularly those related to the immune system. The immune system plays a critical role in embryo implantation, placental development, and the maintenance of pregnancy. However, improper regulation of immune responses can lead to implantation failure, pregnancy loss, or complications during IVF. This section will explore key immunological challenges, including immune rejection and inflammation, which affect IVF outcomes.

**1. Immune Rejection of the Embryo**

One of the primary immunological challenges in IVF is the risk of immune rejection of the embryo. The embryo is a semi-allograft, meaning it carries paternal antigens that are foreign to the maternal immune system. The maternal immune system must, therefore, develop mechanisms to tolerate these foreign antigens while still providing immune protection against potential pathogens.

**Failure of Immune Tolerance:** During a successful pregnancy, the maternal immune system adapts to tolerate the semi-allogeneic embryo through the action of regulatory T cells (Tregs) and the expression of non-classical HLA molecules, such as HLA-G, by the trophoblast. However, in some women undergoing IVF, this immune tolerance may not be properly established, leading to the immune system attacking and rejecting the embryo, contributing to implantation failure [5].

**Elevated NK Cell Activity:** Natural killer (NK) cells play a dual role in pregnancy: they are essential for trophoblast invasion and vascular remodeling but can also become overactive and cytotoxic, leading to embryo rejection. Elevated NK cell activity has been associated with recurrent implantation failure (RIF) and miscarriage. In IVF patients, elevated levels of cytotoxic NK cells may interfere with the establishment of immune tolerance, leading to the rejection of the transferred embryo [6].

**2. Inflammation and Cytokine Imbalance**

Inflammation, mediated by an imbalance of cytokines, is another significant challenge in IVF. Successful implantation requires a balance between pro-inflammatory and anti-inflammatory cytokines. Dysregulation of this balance can lead to a hostile uterine environment, preventing embryo implantation.

**Pro-Inflammatory Cytokines (Th1 Response):** Cytokines such as tumor necrosis factor-alpha (TNF-α), interleukin-6 (IL-6), and interferon-gamma (IFN-γ) are part of the pro-inflammatory Th1 response. While a mild inflammatory environment is necessary for embryo implantation, excessive production of these cytokines can lead to chronic inflammation, creating a hostile uterine environment. Elevated levels of TNF-α and IFN-γ have been associated with implantation failure and recurrent miscarriage in IVF patients [4].

**Anti-Inflammatory Cytokines (Th2 Response):** Anti-inflammatory cytokines, such as interleukin-10 (IL-10) and interleukin-4 (IL-4), are part of the Th2 response, which promotes immune tolerance and a favorable environment for implantation. A deficiency in Th2 cytokine activity or a failure to shift from a pro-inflammatory (Th1) to an anti-inflammatory (Th2) environment can hinder successful implantation [5].

**Imbalance in Th1/Th2 Cytokine Ratios:** The delicate balance between Th1 and Th2 cytokines is crucial for successful implantation and pregnancy. An abnormal shift towards a Th1-dominant response has been implicated in pregnancy loss and poor IVF outcomes. Conversely, a Th2-dominant environment is more conducive to embryo implantation and the continuation of pregnancy [4].

**3. Autoimmune Disorders and Reproductive Failure**

Autoimmune disorders are another challenge for women undergoing IVF. Autoimmune conditions can lead to the production of antibodies that attack the body's own tissues, including those of the reproductive system. These antibodies can interfere with embryo implantation and increase the risk of miscarriage.

**Antiphospholipid Syndrome (APS):** APS is an autoimmune disorder characterized by the presence of antiphospholipid antibodies, which can lead to blood clot formation and placental insufficiency. Women with APS have a higher risk of recurrent miscarriage and implantation failure, and they may require anticoagulation therapy to improve IVF outcomes [8].

**Antinuclear Antibodies (ANA):** ANA are autoantibodies that target the body's cells and tissues. Elevated ANA levels have been associated with implantation failure and recurrent miscarriage, particularly in women undergoing IVF. These antibodies may interfere with the maternal immune system's ability to tolerate the embryo [7]

**4. Endometrial Receptivity and Immune Dysregulation**

For successful embryo implantation, the endometrium must be receptive at the time of embryo transfer. Immune dysregulation in the uterine environment can disrupt this receptivity, preventing the embryo from implanting properly.

**Endometrial NK Cells:** uNK cells play a role in the remodeling of uterine blood vessels during early pregnancy. However, increased numbers of cytotoxic NK cells in the endometrium have been associated with implantation failure. In IVF, women with elevated uNK cell activity may experience repeated implantation failure due to an overly aggressive immune response in the endometrium [6].

**Chronic Endometritis:** Chronic endometritis, characterized by persistent inflammation of the endometrium, can negatively affect endometrial receptivity. Chronic inflammation, often caused by bacterial infections or immune dysregulation, has been linked to poor IVF outcomes, including implantation failure [7].

**5. Immunosuppressive Therapies and Their Risks**

While immunosuppressive therapies, such as corticosteroids, intravenous immunoglobulin (IVIG), and intralipid infusions, are often used to modulate immune responses in IVF, they come with potential risks.

**Excessive Immunosuppression:** Overuse of immunosuppressive therapies can compromise the maternal immune system's ability to defend against infections. This can lead to an increased risk of infections during pregnancy and may negatively impact both maternal and fetal health [9].

**Tailored Treatment:** Immunosuppressive therapies must be carefully tailored to each patient's unique immunological profile. Over-suppression of immune responses may lead to complications, while inadequate suppression may result in persistent immune rejection of the embryo [7].

3.5 Immunological Prospects in IVF

The immune system plays a critical role in the success of in vitro fertilization (IVF), particularly in embryo implantation and the early stages of pregnancy. During this process, a delicate balance is required between maternal immune tolerance and protection from pathogens. Key immune cells and cytokines orchestrate this balance, promoting a conducive environment for implantation and placental Development. Understandingthe role of these immune factors can offer potential strategies for enhancing IVF outcomes.

**1. Immune Cells in IVF**

Several types of immune cells contribute to a successful IVF outcome by promoting immune tolerance and preventing excessive inflammatory responses:

**a. Regulatory T Cells (Tregs)**

Tregs are critical in establishing immune tolerance during pregnancy, helping prevent the maternal immune system from attacking the embryo, which contains paternal antigens. Tregs suppress the activity of other immune cells which could lead to embryo rejection. Studies have demonstrated that increased Treg populations are associated with successful pregnancies, both in natural conception and IVF [6].

**b. Natural Killer (NK) Cells**

Uterine natural killer (uNK) cells are the most abundant immune cells in the endometrium and play a crucial role in the early stages of pregnancy. These cells help regulate trophoblast invasion and vascular remodeling, which are essential for the development of a healthy placenta.

However, excessive activity of NK cells, particularly cytotoxic NK cells, has been associated with recurrent implantation failure [6]. This dual role makes NK cells both vital for pregnancy success and a potential target for intervention in cases of IVF failure.

**c. Dendritic Cells**

Dendritic cells (DCs) act as antigen-presenting cells that can influence the immune response at the maternal-fetal interface. The proper function of DCs is important for promoting tolerance to the embryo. Some evidence suggests that the maturation and migration of DCs in the endometrium are involved in regulating the local immune environment, which can affect implantation success [10].

**2. Cytokines and Immune Modulation**

Cytokines, small proteins that mediate immune responses, are critical to regulating the balance between pro-inflammatory and anti-inflammatory signals at the maternal-fetal interface. Cytokine balance is crucial for creating an environment conducive to embryo implantation and placental development.

**a. Th1/Th2 Cytokine Balance**

Successful implantation and pregnancy depend on a balance between pro-inflammatory Th1 cytokines (e.g., IFN-γ, TNF-α) and anti-inflammatory Th2 cytokines (e.g., IL-4, IL-10). Th1 cytokines are associated with inflammation and are typically downregulated during early pregnancy to avoid an excessive immune response that could harm the embryo. On the other hand, Th2 cytokines promote immune tolerance, which is necessary for the embryo to implant and thrive [4].

**b. Interleukin-10 (IL-10)**

IL-10 is a critical anti-inflammatory cytokine that promotes immune tolerance at the maternal-fetal interface. Elevated levels of IL-10 are often found in successful pregnancies, and it plays a protective role by inhibiting the production of pro-inflammatory cytokines and preventing excessive immune responses that could lead to implantation failure [5].

**c. Transforming Growth Factor-Beta (TGF-β)**

TGF-β is another anti-inflammatory cytokine that is crucial for promoting immune tolerance during early pregnancy. It helps regulate the activity of Tregs and supports the establishment of a maternal-fetal interface conducive to embryo implantation. Aberrant expression of TGF-β has been linked to pregnancy complications, including implantation failure [7].

**3. Immunological Interventions**

Given the importance of immune regulation in IVF, several immunotherapies have been explored to enhance success rates, particularly in cases of recurrent implantation failure (RIF):

**Intralipid Therapy:** Intralipid, a fat emulsion, has been investigated for its potential to modulate immune responses by reducing NK cell activity. It is believed to improve the chances of embryo implantation in women with elevated NK cell levels [9].

**Intravenous Immunoglobulin (IVIG):** IVIG has been used in reproductive immunology to treat women with recurrent miscarriages or implantation failures, possibly by modulating the activity of Tregs and NK cells and enhancing immune tolerance [7].

**Corticosteroids:** Low-dose corticosteroids are sometimes used in IVF to suppress harmful immune responses and reduce inflammation, potentially aiding in implantation success by promoting a more favorable uterine environment [8].

The maternal immune system plays a pivotal role in determining the success of IVF. Key immune cells, such as Tregs and NK cells, and cytokines like IL-10 and TGF-β, are essential for creating an environment that supports embryo implantation and early pregnancy. Advances in reproductive immunology, particularly in understanding immune modulation, have opened new avenues for enhancing IVF outcomes through targeted immunotherapies.

**4.0 Future Directions of Immunological Research in IVF**

The role of immunology in in vitro fertilization (IVF) continues to be an area of intense study, as the success of IVF is strongly influenced by the immune system's regulation of implantation and pregnancy. Despite advances in reproductive immunology, several unresolved challenges remain, including recurrent implantation failure (RIF), pregnancy loss, and unexplained infertility. Future research efforts are focused on unraveling the complex interactions between the maternal immune system and the developing embryo, as well as developing immunomodulatory therapies that can enhance IVF outcomes. This section outlines several promising avenues for future research in this field.

**1. Personalized Immunotherapy for IVF Patients**

As the understanding of immune profiles in women undergoing IVF deepens, there is growing interest in developing personalized immunotherapy tailored to individual patients. Current immunomodulatory treatments, such as intravenous immunoglobulin (IVIG), intralipids, and corticosteroids, are often used empirically in women with recurrent implantation failure or miscarriage. However, the success of these therapies remains inconsistent due to a lack of precise diagnostic markers.

**Personalized Immune Profiling:** Future research may focus on developing more sophisticated diagnostic tools to identify immune abnormalities in women undergoing IVF. This could involve comprehensive immune profiling, including detailed analysis of T cell subsets, natural killer (NK) cell activity, and cytokine levels, to create tailored treatment plans that address the specific immune imbalances of individual patients [7].

**Targeted Immunotherapies:** In the future, treatments such as monoclonal antibodies or small-molecule inhibitors may be developed to precisely modulate specific immune pathways. For instance, therapies targeting overactive NK cells or pro-inflammatory cytokines, such as TNF-α or IFN-γ, could improve implantation rates in women with elevated immune activity [6].

**2. Role of the Microbiome in Reproductive Immunology**

Emerging evidence suggests that the uterine and vaginal microbiome may play a significant role in regulating immune responses during pregnancy and influencing IVF outcomes. The composition of the microbiome has been linked to both immune activation and tolerance in the reproductive tract, making it a promising target for future research.

**Microbiome-Immune Interaction:** Future studies will likely focus on understanding how the microbiome interacts with the maternal immune system during IVF. For example, certain bacterial species may promote immune tolerance and reduce inflammation, while others may trigger immune responses that hinder implantation (Moreno et al., 2016). Modulating the microbiome through probiotics or targeted antibiotics could be explored as a strategy to optimize the immune environment for embryo implantation.

**Microbiome profiling as a Diagnostic Tool:** Profiling the uterine and vaginal microbiome could become a routine part of infertility diagnostics. Personalized interventions to restore a healthy microbiome may enhance endometrial receptivity and improve IVF outcomes [11].

### 3. Epigenetic Regulation of Immune Responses in IVF

Epigenetics, which refers to changes in gene expression without alterations in the DNA sequence, has gained increasing attention as a potential factor in reproductive success. Epigenetic mechanisms, such as DNA methylation and histone modifications, can influence the function of immune cells, including Tregs and NK cells, at the maternal-fetal interface.

**Epigenetic Modulation of Immune Cells:** Future research may explore how epigenetic changes in immune cells affect their role in pregnancy. For example, alterations in the epigenetic landscape of Tregs or NK cells could impair their ability to promote immune tolerance during embryo implantation [12]. Targeting these epigenetic modifications with drugs that restore normal gene expression could represent a novel therapeutic strategy for improving IVF outcomes.

**Environmental and Lifestyle Factors:** Environmental factors such as stress, diet, and exposure to toxins can influence epigenetic regulation. Future studies may investigate how these factors impact the immune system and fertility, potentially leading to lifestyle recommendations or interventions that promote a favorable immune environment for IVF [12].

### 4. Immunomodulatory Agents for Enhancing Endometrial Receptivity

One of the most critical factors influencing IVF success is endometrial receptivity, which is partially regulated by immune cells such as Tregs, NK cells, and macrophages. Research into immunomodulatory agents that specifically target the uterine immune environment could lead to improved therapies for women with implantation failure.

**Cytokine Therapies:** Cytokine therapies that enhance endometrial receptivity by modulating immune responses at the maternal-fetal interface are a potential future direction. For example, interleukin-10 (IL-10), a key anti-inflammatory cytokine, has been shown to promote immune tolerance during implantation. Future research could explore the use of cytokine-based therapies to improve implantation rates [8].

**Stem Cell Therapy:** There is growing interest in using mesenchymal stem cells (MSCs) to enhance endometrial receptivity by modulating local immune responses. MSCs have been shown to suppress inflammation and promote tissue repair, making them a promising option for women with poor endometrial receptivity or chronicendometritis [13].

### 4.2. Immunogenetic Screening and Genetic Counseling

Advances in immunogenetics offer new opportunities to predict and prevent immune-related complications in IVF. Genetic variants in immune-related genes, such as those coding for human leukocyte antigens (HLAs), cytokines, and other immune regulators, can influence the maternal immune response to the embryo.

**Immunogenetic Testing:** In the future, immunogenetic testing could become a routine part of fertility assessments, allowing clinicians to identify genetic predispositions to immune dysregulation, such as abnormal NK cell activity or cytokine production. This information could be used to customize treatment plans, including the use of immunomodulatory therapies [7].

**Gene Editing and CRISPR:** In the longer term, gene-editing technologies like CRISPR could be explored as tools to correct immune-related genetic defects that contribute to infertility or pregnancy loss. While these technologies are still in their infancy, they offer the potential for precise intervention in immune pathways that are critical for IVF success [12].

### 5.CONCLUSION

Immunological challenges, including immune rejection, inflammation, cytokine imbalances, and autoimmune disorders, significantly affect the success of IVF. The delicate balance between immune tolerance and immune defense is critical for embryo implantation and early pregnancy. Advances in reproductive immunology, particularly in the development of targeted immunotherapies, offer potential solutions to overcome these challenges, but careful management is essential to avoid unintended consequences. Continued research into the intricate immune mechanisms governing implantation and pregnancy will be crucial in developing targeted treatments to overcome the immunological challenges associated with IVF.

**Declaration of conflict of interest:** The authors declare no conflict of interest.

**Funding:** None

**Data availability statement:** Data are available upon request from the first author or any of the other authors.

### REFERENCES

[1]Sacks, G. More than a barrier: Immune system's role in successful implantation in IVF. Fertility and Sterility, 2014; 102 (2): 299-305.

[2]Kushnir, V. A., Barad, D. H., Albertini, D. F., &Gleicher, N. Oocyte cryopreservation: A realistic and practical option for fertility preservation. Reproductive Biology and Endocrinology, 2016

[3]Society for Assisted Reproductive Technology 2021. IVF success rates. Retrieved from https://www.sart.org

[4]Saito, S., Nakashima, A., Shima, T., & Ito, M. Th1/Th2/Th17/Th22 and regulatory T-cell paradigm in pregnancy. American Journal of Reproductive Immunology, 2015; 63(6): 601-610.

[5]Robertson, S. A., &Moldenhauer, L. M. Immunological determinants of implantation success. Human Reproduction Update, 2014; 20(1): 86-104.

[6]Lachapelle, M. H., Miron, P., Hemmings, R., & Roy, D. C. Endometrial NK cells in patients with recurrent miscarriage and implantation failure: An immunological paradox. Human Reproduction, 2017; 12(3): 393–397.

[7]Kwak-Kim, J., & Gilman-Sachs, A. Reproductive immunology: Implications for pregnancy and IVF outcomes. Current Opinion in Obstetrics and Gynecology, 2018; 30(4): 239-246.

[8]Rai, R., & Regan, L. Antiphospholipid antibodies and reproductive failure. British Medical Journal, 2017; 310(6979): 704-707.

[9]Gibson, K. S., Dietz, L. D., Little, B. B., Rinehart, J. S., & Joiner, J. S. Effects of intralipid infusion on natural killer cell cytotoxicity and pregnancy outcomes in women undergoing in vitro fertilization. Fertility and Sterility, 2015; 103(2): 529-534.

[10]Blois, S. M., Barrientos, G., Garcia, M. G., & Orsal, A. S. Dendritic cell function during pregnancy: Ample prospects for diverse therapeutic approaches. American Journal of Reproductive Immunology, 2011; 65(1): 114-123.

[11]Moreno, I., Codoner, F. M., Vilella, F., Valbuena, D., Martinez-Blanch, J. F., Jimenez-Almazan, J., Simon, C. Evidence that the endometrial microbiota has an effect on implantation success or failure. American Journal of Obstetrics and Gynecology, 2016; 215(6): 684-703.

[12]Jansson, M. D., Damasco, C., &Demaria, M. Epigenetic regulation of immune responses during pregnancy and its implications for reproductive health. Trends in Immunology, 2019; 40(9): 760-774.

[13]Tan, J., Li, P., Wang, Q., & Wang, L. Mesenchymal stem cells as therapeutic agents for infertility and reproductive disorders. Stem Cell Research & Therapy, 2020; 11(1): 200.

© Copyright 2025 BioMedSciDirect Publications IJBMR -ISSN: 0976:6685.