



Contents lists available at BioMedSciDirect Publications

International Journal of Biological & Medical Research

Journal homepage: www.biomedscidirect.com

Original Article

THE BIOACTIVITY OF HEXADECANOIC ACID COMPOUND ISOLATED FROM HYDROID *Aglaophenia cupressina* LAMOUREOUX AS ANTIBACTERIAL AGENT AGAINST *Salmonella typhi*

Eva Johannes^a, Magdalena Litaay^b, Syahribulan^c^{a,b,c} Department of Biology, Mathematic and Natural Sciences Faculty, Hasanuddin University, Jl. Perintis Kemerdekaan Km 10 Tamalanrea Makassar 90245 Makassar

ARTICLE INFO

Keywords:

Bioactivity,
isolates,
Hydroid,
Aglaophenia,
antibacteria,
Salmonella.

ABSTRACT

Background & Objectives: The new, more effective antibiotic sources are urgently needed now to overcome the multi-drug resistance (MDR) in the treatment of infectious diseases caused by pathogenic microorganisms. This study was aimed to find out the bioactivity of the hexadecanoic acid isolated from hydroid *Aglaopheniacupressina*Lamoureaux with inhibition mechanisms or lethal to *Salmonella typhi* that frequently contaminate food materials. **Methods:** This study was an experimental study with the following treatment steps: Isolation and characterization of the compounds from hydroid *A. cupressina*Lamoureaux by chromatography. Hexadecanoic acids and the isolates from hydroid *A. cupressina*Lamoureaux were then tested against *S. typhi* at various concentrations (10 ppm, 20 ppm, and 30 ppm). **Result:** Study findings indicated that: (1) hexadecanoic acid at the concentration of 30 ppm is bactericidal against *S. typhi*. **Interpretation & Conclusion:** From the study findings, it can be concluded that hexadecanoic acid (30 ppm) has antibacterial activity by damaging the cells wall of *S. typhi*.

© Copyright 2010 BioMedSciDirect Publications IJBMR - ISSN: 0976:6685. All rights reserved.

1. Introduction

The main problem in the treatment of infectious diseases caused by pathogenic microorganisms is the emergence of the multi-drug resistant (MDR) pathogens. This issue has become an important focus of researches on new bioactive compounds. According to Murniasih, the development of new medicines derived from marine biota is attracting researchers attention due to the very high marine biodiversity and the uniqueness of the produced secondary metabolites structure^{1,2}. This can be achieved by isolating bioactive compounds from marine natural materials, which are safe for health while capable of inactivating bacteria and fungi that contaminate food materials. These active substances accumulate in high number in some marine invertebrates including sponges, tunicates, mollusks, and cnidaria³.

The hydroid *Aglaophenia cupressina* Lamoureaux is an invertebrate from cnidaria phylum that lives by attaching itself to sponges. In addition to alkaloid, diterpene, tridentatol A, and prostaglandin, it also contains histamines, protein and histamine liberator in its nematocyst⁴ also confirmed that rough extract of *Aglaophenia* sp (0.05%) is capable of inhibiting the growth of *Fusariumoxysporum* f. Sp. *Vanilae*. It is assumed that there are remaining many other bioactive compounds from secondary metabolites of hydroid *A. cupressina*L. that can be utilized as pesticide, antifungal and antibacterial. According to⁵, secondary metabolite compounds can be treated as lead compound in agricultural and medicinal industries.

This study was a continuation of a previous study on isolation and characterization of secondary metabolites from hydroid *A. cupressina*Lamoureaux as antimicrobial precursor⁶. From the discovered compounds, one of them was hexadecanoic acid with antibacterial activity in *Escherichia coli* and have been published⁷. This study was intended to know the ability of hexadecanoic acid and its mechanism in inhibiting or inactivating the Gram-negative bacteria growth such as *Salmonella typhi* which is a pathogenic bacterium that frequently contaminate food materials and result in intoxication. According Schneider et al. (2008), *S. typhi* is not inactivated by cells after consumption, but multiplying itself in cells, and then out of the cells to blood circulation, spreading to entire body that result in systemic infections.

Material and Methods

This study was conducted in Microbiological Laboratory of Hasanuddin University from May to September 2014 with experimental method.

Material

Material used in this research such as hydroid *Aglaophenia cupressina* Lamoureaux, hexadecanoic acid compound, pure culture (*S. typhi*), dimethyl sulphoxide (DMSO) (E Merck), chloramphenicol (PT Alpha), physiological NaCl 0.9%, glucose nutrient agar (GNA) medium, nutrient agar (NA) medium, nutrient broth (NB) medium, Muller Hinton agar (MHA) medium (oxid).

Procedure

Extraction, Partition and Isolation

* Corresponding Author :

Tlp/Fax: 081944210934.

Email : evajohannes@gmail.com

©Copyright 2010 BioMedSciDirect Publications. All rights reserved.

The collected samples were cleaned from their substrate, washed, and macerated with methanol for 24 hours three times, filtered and evaporated to obtain thick macerates. The macerates were then partitioned (liquid-liquid 1:1) with n-hexane. Filtrate or n-hexane layer was evaporated to obtain thick extract of n-hexane and continued with vacuum chromatography column, and pressured chromatography column to obtain pure compounds. The obtained pure compounds were then analyzed by thin layer chromatography and melting point measurement, using the reference hexadecanoic acid⁷.

Antibacterial Activity Test using Agar Diffusion Method

20 mL of Sterile Muller Hinton agar (MHA) was poured aseptically into petri disc and left to solidify as base layer. After that, the suspension of test bacteria was added 1 mL each into 10 mL of medium above the base layer and left to semi-solid as seeding layer. Six parts with inner diameter of 5 mm, outer diameter of 8 mm, height of 10 mm, were put aseptically with sterile pincer on medium surface at interval of 2-3 cm from petri disc edges, and then stored at room temperature. Each of the parts was added with 0.25 mL hexadecanoic acid and β -cytosterol isolated from hydroid *A. cupressina* Lamoureux at concentrations of 10 ppm, 20 ppm, and 30 ppm. Chloramphenicol as positive control and DMSO as negative control, 0.25 mL each, and then incubated at 37°C for 24 hours and 48 hours. Observations were performed by measuring the inhibitory diameter of bacterial growth around the parts by using calliper to observe the ability of the compound in inhibiting the test bacterial growth. The inhibitory ability measurement results at 24 hours and 48 hours were tabulated and analysed.

Results And Discussion

Analysis of Hexadecanoic Acid Structure

Yellowish white crystal weighting 50 mg with melting point of 430 – 440 C. The compound was fluorescent under UV and blue in colour, but it was not visible by TLC colour test. UV spectrum (CH₃OH) indicated maximum absorption at λ_{max} of 212 (164) nm; IR spectrum (KBr) ν_{max} of 3472 cm⁻¹ (OH), 2921, 2855 cm⁻¹ (aliphatic C-H), 2672 cm⁻¹ (aliphatic C-H), 1707 cm⁻¹ (C=O), 1466 cm⁻¹ and 1415 cm⁻¹ (CH₂ and CH₃), 1299 cm⁻¹ (C-O). NMR spectrophotometer analysis includes ¹H-NMR, ¹³C-NMR, and HMBC as indicated in Figure (1) and Table (1).

Hexadecanoic acid compound had 16 carbons, which were identified by signals appeared as δ C 180.64 signals with the most distant chemical shift indicated carbon at carboxyl group, signals at δ C 34.30 and 32.12 indicated C-2 and C-14. There are 6 carbons in symmetrical position, namely: C-7, C-8, C-9, C-10, and C-11. It provided one signal with high intensity, namely δ C 29.89. In addition, there were very near signals at δ C 29, 89; 29, 79; 29, 56; 29, 44; 29, 25, each of them represented C-4; C-5; C-13; C-6 and C-12. Signals at δ C 24.85 ; 22.88 and 14.31 represented C-13, C-15 and C-16, respectively.

Analysis of ¹H-NMR spectrum indicated the following signals: δ H 2,33 (2H,t, J = 7.35) representing 2 proton at C-2, signal at δ H 1.62 (2H,m, J =7.35) representing 2 protons at C-3, signal at δ H 1.24-1.28 (20H,m) indicated 20 hydrogen at C-4; C-5; C-6; C-7; C-8; C-9; C-10; C-11; C-12 and C-13, signal δ H 1,28-1,29 (2H, m) represented 2 protons at C-14, signal at δ H = 1,31-1,33 (2H, m) representing 2 protons at C-15, signal

with δ H = 0,87 (3H, t, J =6.70) represented 3 protons at C-19, signal at 3.75 (1H, s) represented proton at hydroxyl group. The number of hydrogen in the compound were 32.

The compound structure elucidation was supported by HMBC spectrum that indicated distant correlation between proton signals and carbon signals as follow; proton signal δ H 2,33 (H-2) correlated distantly to carbon signals δ C 180,64 (C-1) and 24,85 (C-3), proton signal δ H 1,28-1,29 (H-14) correlated to signal δ C 22,88 (C-15). Proton signals δ H 1,31-1,33 (H-15) correlated to carbon signals δ C 32,12 (C-14) and δ C 14,31 (C-16) and the correlation between proton signal δ H 0,87 (H-16) and carbon signals δ C 32,12 (C-14) and δ C 32,12 (C-14) and δ C 22, 88 (C-15).

Antibacterial Activity by Inhibitory Diameter Measurement

Inhibitory diameter measurement in *Salmonella typhi*

The results of inhibitory diameter measurement indicated that antibacterial bioactivity of hexadecanoic acid against *S. typhi* at the concentrations of 10ppm and 20ppm for incubation at 24 hours did not subject to inhibitory zone change compared to incubation at 48 hours. In contrast, at concentration of 30ppm the inhibitory zone at 24 hours was 16.00 ppm, but at incubation period of 48 hours the inhibitory zone increased to 17.00 ppm. This indicated that the hexadecanoic acid at concentration of 10 ppm and 20 ppm was bacteriostatic, whereas at the concentration of 30 ppm it was bactericidal. This means that at concentration of 30ppm, hexadecanoic acid not only inhibit the *S. typhi* growth but also lethal.

According to Madigam et al. (2012), the bacteriostatic properties can change into bactericidal when the concentration of the compound gets bigger. The mechanism of bactericidal compound affects the bacterial growth by damaging and breaking the cells wall so the bacteria cannot stand with external effect, or disturbing the bacterial cell membrane wholeness, that the active substances or metabolites exchange in and out of the cells will be disturbed¹⁰.

Chloramphenicol used as positive control was an aminoglycoside antibiotic, which was bacteriostatic that did not kill the bacteria but inhibit the synthesis of the proteins that are urgently needed in multiplication and division of bacterial cells¹¹. Whereas the DMSO used as negative control did not have antibacterial properties, used to dissolve and stabilize the intracellular extract to be tested¹².

Mechanisms of Bacterial Cells Wall Chemical Reaction to Hexadecanoic Acid

Figure 6 shows the esterification reaction between hexadecanoic acid and hydroxyl group of lipopolysaccharide that compose the cells wall of gram-negative bacteria. The reaction resulted in alteration in lipopolysaccharide membrane structure into asymmetric. This disturbs the balance in the membrane lipid structure, so the integration and flexibility of the cells membrane are perturbed, which according to Kim et al. (1995) the reaction between membrane active components and antibacterial compound results in changes in membrane compositions followed by cells swelling and then cytoplasm membrane damage, distended and lysed. According to Radiati (2002), the OH group of hexadecanoic acid can be toxic to cell protoplasm, infiltrating and damaging the cells wall and denaturize the protein in the cytoplasm and form hydrogen bond on the active site of enzyme.

Table 1. ¹H-NMR and ¹³C-NMR Data of Hexadecanoic Acid Compound⁷

No.	¹³ C-NMR δ _c (ppm)	¹ H-NMR δ _H : ppm (multiplicity, J in Hz)	HMBC
1	180,64	-	-
2	34,3	2.33 (2H, t, J = 7.35)	1,3
3	24,85	1.62 (2H, m, J = 7.35)	1,2
4-	29,89; 29,79; 29,56; 29,44;		
13	29,25	1.24 - 1.28 (20 H, m)	-
14	32,12	1.28 - 1.29 (2H, m)	15
15	22,88	1.31 - 1.33 (2H, m)	14,16
16	14,31	0.87 (3H, t, J = 6.70)	14,15
OH	-	3.75 (1H, s)	-

Table 2. Antibacterial bioactivity of hexadecanoic acid against Salmonella typhi at the concentrations of 10ppm and 20ppm for incubation at 24 and 48 hours

Treatment	Inhibitory Zone Diameter (mm)	
	24 hour	48 hour
A = Hexadecanoic acid at 10ppm	18,00	18,00
B = Hexadecanoic acid at 20ppm	18,50	18,50
C = Hexadecanoic acid at 30ppm	16,00	17,00
D = Negative Control (DMSO)	0,00	0,00
E=Positive Control (Chloramphenicol)	12,00	15,00

Source; processed data 2014

Fig 1. NMR Spectrometer Analysis




Fig. 2. Salmonella typhi, incubation at 24 hours



Fig 3. Salmonella typhi, incubation at 48 hours





Fig 4. Bacterial Cells Wall Reaction to Hexadecanoic Acid



Conclusion

Hexadecanoic acid at concentration of 30 ppm is bactericidal against Salmonella typhi by damaging its cell membrane structure.

Acknowledgement

The authors thank to Directorate of Higher Education for its financial support in this study through BOPTN UNHAS 2014.

References

1. Muniarsih, T., 2003. Secondary Metabolites from Sponge as Medicinal Materials. J. Oseana, 3:27-33.
2. Joana, Luisa Peixe, Newton C.M. Gomes, Ricardo Calado, 2011. Cnidarians as a Source of New Marine Bioactive Compounds—An Overview of the Last Decade and Future Steps for Bioprospecting. (www.mdpi.com/journal/marinedrugs). Rev., accessed 20 July 2012).
3. Xue Song, Hai Tao Zang, Pei Chun Wu, Wei Zhang, Quan Yuan., 2004. Journal of Experimental Marine Biology and Ecology. 298., 71-78. www.elsevier.com/locate/jembe.
4. Suada I Ketutdan Ni WayanSuniti, 2010. Suppression ability of Crude Extract Derived from Marine Biota Against Fusarium oxysporumf.sp.vanillae. Jurnal Biologi, XIV (1): 7-10.

5. Rachmaniar, R., 2003. Anticancer Swinholide A from Sponge Theonella Swinhoei. *J. Bahan Alam Indonesia*. Vol 2 No 4, 122.
6. Johannes E., 2008. Isolation, Characteristic and Secondary Metabolite Bioactivity Test on Hydroid *Aglaophenia cupressina* Lamoureaux as Antimicrobial Basic Material. Thesis. Postgraduate Program, Hasanuddin University.
7. Johannes E. Hanapi Usman, Mariyati Bilang., 2013. Effectiveness of Extracted Antibacterial Compound from Hydroid *Aglaophenia cupressina* Lamoureaux Against Bacterial Cell of *Escherichia coli*. *International Journal of Scientific and Technology Research* Vol.2. Issue 2. ISSN : 2277-8616
8. Schneider, R. K. and Rence M. Goodrich, 2008. Dealing With Foodborne Illness: Typhoid Fever. *Salmonella typhi*. *J. Solution for Your Life*. <http://edis.ifas.ufl.edu/fs.125>, (accessed 20 July 2014).
9. Madigam M. T, Martinko J. M, Dunlap, P.V, Clark, D. P., 2012. *Biology of Microorganism* 13th ed. San Francisco : Pearson. p. 34.
10. Pelczar MJ, Chan ECS, 2006. *Microbiology Basics 2*. Penerbit Universitas Indonesia. Jakarta.
11. Bhat V., 2008. Classification of Antibiotic. Medical notebook. From <http://pre-pg.blogspot.com/2007/03/classification-ofantibiotics.html>.
12. Pahusip Adolf, Sedarwati Yasin, Yenni Elisabeth, 2003. A Study on Extraction Method of Andaliman (*Zanthoxylum acanthopodium* DC) Against Pathogenic and Food-Damaging Microbes. *Jurnal Ilmu dan Teknologi Pangan*. Vol. 1 No. 1.
13. Kim JM, Marshal MR, Cornell JA, Boston JF, Wei CL., 1995. Antibacterial Activity of Carvacrol, Citral and geraniols against *Salmonella typhimurium* Culture Medium and Fish Cubes. *J. Food Sci* 60 (6): 1365-1368.
14. Radiati LE. 2002. Inhibition Mechanism of Enteropathogenic Bacterial Virulence by Ginger Rhizome (*Zingiber officinale* roscoe) [Dissertation] Bogor. Postgraduate Program, Bogor Agricultural Institute.