

**Original article****Antibacterial activity of *Daniellia oliveri*, *Lophira alata*, and *Lannea welwitschii* against plasmid-carrying enteric pathogens from Lagos, Nigeria**KO Akinyemi<sup>a</sup>\*CO Fakorede<sup>a</sup>, BA Iwalokun<sup>b</sup>AO B Oyefolu<sup>a</sup><sup>a</sup>Department of Microbiology, Lagos State University, Ojo P.M.B 1087, Apapa, Lagos, Nigeria<sup>b</sup>Biochemistry Unit and Malaria research Laboratory, Nigerian Institute of Medical Research, Yaba, Lagos, Nigeria

## ARTICLE INFO

**Keywords:**Plant,  
extract,  
MDR,  
pathogens,  
plasmid,  
curing,  
photochemical

## ABSTRACT

**Aim:** Three medicinal plants *Daniellia oliveri*, *Lophira alata* and *Lannea welwitschii* were tested for in vitro antibacterial activity and their plasmid eviction potentials on enteric pathogens. **Method:** Fresh plant materials collected from herbs seller were sun dried and grounded to fine powder. Water and methanol extracts of these plants were obtained by standard methods. Enteric bacterial agents from clinical samples were used. In-vitro antimicrobial susceptibility testing of the bacterial isolates was carried out using commercial antibiotics and the crude plant extracts by standard procedures. Phytochemical screening of the plant extracts was carried by the standard procedure. Plasmid DNA extraction was performed using the alkaline lyses method. **Result:** The crude methanol extracts of all the plants were effective on MDR strains of *Salmonella enterica* serova Typhi, *S. Typhimurium*, *Escherichia coli*, *Shigella* spp and *Klebsiella* spp used in this study MIC and MBC values of 0.625 to 10 mcg/ml and 5.0 to 20 mcg/ml respectively were recorded for *L. welwitschii*, 10 to 20 mcg/ml and 20 to 40 mcg/ml for *D. oliveri*, and 10 to 20 mcg/ml and 20 to 40 mcg/ml were recorded for *L. alata* respectively. The plasmid eviction potentials of the crude plants extracts indicated that only crude extract of *L. welwitschii* exerted anti-plasmidic effect on *Shigella* spp, *Klebsiella* spp, *S. Typhimurium*, and *S. Typhi*, as over 60% of these bacterial strains loss their plasmid DNA. **Conclusion:** This study revealed that methanol extracts of the three medicinal plants were active against enteric bacterial pathogens. The extracts of *L. alata* and *D. oliveri* were not active against MDR pathogens thus; their continuous usage should be discouraged. The methanol crude extract of *L. welwitschii* exerted antiplasmidic effect on MDR enteric pathogens. This finding suggests possibility of a new approach in drug formulation using anti-plasmidic plant extracts for effective management of plasmid encoded drug resistant enteric bacteria-associated diseases

© Copyright 2010 BioMedSciDirect Publications IJBMR - ISSN: 0976:6685. All rights reserved.

**1. Introduction**

Enteric infection is an increasingly recognized array of bacterial, parasitic and viral pathogens which can profoundly disrupt intestinal function with or without causing overt dehydrating diarrhoea. Some of the organisms implicated in enteric infection include *Salmonella* spp., *Shigella* spp., *Proteus* spp., *Klebsiella* spp., *Staphylococcus aureus* and *Escherichia coli*. Enteric pathogen has been reported as the major causative agent of sporadic and endemic diarrhoea both in children and in adult [1]. Drugs like aminoglycosides, cephalosporins, and fluoroquinolone have antibacterial effects against enteric pathogens [2]. The evolutionary pressure from the use of

antibiotics has played roles in the development and the spread of multidrug resistance among different bacteria species. Also, extra-chromosomal genetic elements (plasmids) harboured by some bacterial agents had been responsible for increased resistant development either horizontally or vertically [2]. The use of medicinal plants and herbs in curing diseases has always been part of human culture and has transcended all social, cultural, economic, religious and other barriers created by man. It has been established that the plants which naturally synthesis and accumulate some secondary metabolites possess medicinal properties [3].

In Nigeria many hundred of plants are used in traditional medicine as treatment for bacterial infections, some of these plants have been subjected to an in-vitro screening but the efficacy of such herbal medicine has seldom been rigorously tested in controlled clinical trials. Plants have been major source of medicine and plants secondary metabolite has been attributed for most plant therapeutic

\* Corresponding Author : **Akinyemi K.O (PhD)**

Department of Microbiology, Lagos State University

PMB 1087, Apapa, Lagos, Nigeria.

Phone: +2348029088676

E-mail: [akinyemiko@yahoo.com](mailto:akinyemiko@yahoo.com)

activities [4-]. *Daniellia oliveri* is of medicinal important value being potent in treating gastrointestinal ailment [6], as antibioficient agent in pregnancy, pain killer, skin, mucosa and sedative [7] for the cure of rheumatism pain [8] and active as an antimicrobial agent [9,10]. In traditional African medicine, *Lannea welwitschii* is used for swellings, oedema, gout, haemorrhoids and emesis [11] antidiarrheal [12] and antipurgative [13]. While in South West Nigeria *Lophira alata* is use for the treatment of typhoid fever [13] and for the treatment of fever, cough, jaundice and gastrointestinal disorder [14]. Therefore a need to search and develop newer, safer, effective and cheaper antimicrobial agent from natural products such as medicinal plants that will compliment synthetic antimicrobial drugs against multiple drug resistant pathogens is paramount importance. This study was undertaken with a view to determining the antimicrobial activity of three medicinal plants and their plasmid curing ability on plasmid-mediated multiple drug resistant enteric bacterial pathogens.

## Materials and methods

### Plant materials

Fresh plant materials of *Lannea welwitschii*, *Daniellia oliveri* and *Lophira alata* were collected from herbs seller in Itoku market, Abeokuta south local government Area of Ogun state Nigeria. The identification and authentication of these plants were carried out by Mr. OK Oluwa and voucher specimens' numbers were deposited in the herbarium of the Botany department of the Lagos State University.

### Extraction

Stem bark of *Daniellia oliveri* (A) Rolfe. Hutch & Dalz, *Lannea welwitschii* (B) Hiern, and *Lophira alata* (C) Bank ex f. Gaertn.f were sun dried for one week and were grounded to fine powder using pestle and mortar and were stored at room temperature in a clean dry air tight bottle for further use.

### Methanol extracts (Cold)

The methanol extractions of these plants were carried as described by Akinyemi et al [14]. In brief twenty grams of the powdered plant materials A, B, and C were weighed and soaked in separate beakers containing 200 ml of 70% methanol each. The beaker was wrapped with aluminium foil and sealed with tape. The mixture was shaken vigorously and left to stand overnight (24 hours) in a refrigerator maintained at 2 to 8°C. The mixture was filtered using a clean muslin cloth. The filtrate was evaporated to dryness in a hot air oven maintained at 40°C for 72 hours. The crude extracts were scrapped and weighed to determine the yield. The crude extracts yield were dispensed in a sterile container and labelled as methanol extracts and designated as AM (*D. oliveri*), BM (*L. welwitschii*), and CM (*L. alata*) respectively.

### Aqueous extract (Cold)

The aqueous extraction method was as described by Akinyemi et al (2005) [15]. In brief twenty grams of the powdered plant materials A, B, and C were weighed and soaked in separate beakers containing 200 ml of distilled water each. The beaker was wrapped with aluminium foil and sealed with tape. The mixture

was shaken vigorously and left to stand overnight (24 hours) in a refrigerator maintained at 2 to 8°C. The mixture was filtered using a clean muslin cloth. The filtrate was evaporated to dryness in a hot air oven maintained at 40°C for 72 hours. The crude extracts were scrapped and weighed to determine the yield. The crude extracts yield were dispensed in a sterile container and labelled as water extracts and designated as AW (*D. oliveri*), BW (*L. welwitschii*), and CW (*L. alata*) respectively.

### Bacterial cultures

The bacteria used in this work were obtained from the Nigerian institute of Medical Research (NIMR), Yaba, Lagos, Nigeria and were all subjected to standard microbiological methods maintained on Tryptone soya agar petri dishes at 2 to 8°C for 24 hrs before use. These bacteria include *Salmonella enterica* serovar Typhi, *Salmonella Typhimurium*, *Escherichia coli* ATCC 25922, *Shigella* spp, and *Klebsiella* spp, and were isolated from stool samples of patients diagnosed of gastro intestinal infections.

### Antimicrobial susceptibility testing

Agar disc diffusion assay: In-vitro susceptibility of the bacterial isolates to the plant extract was tested for lead antimicrobial response by the standard disc diffusion technique using guideline established by the NCCLS 2002 [16]. A standardized Inoculum of  $1.0 \times 10^7$  with 0.5 McFarland standards was used to evenly inoculate the surface of Mueller-Hinton agar plate (Oxoid, UK). Sterile paper discs previously soaked in known concentration (5, 20, and 40 mg/ml) of extracts were carefully placed at different spot on agar plate containing the organisms used and the discs were gently pressed down to ensure contact. The plates were incubated aerobically at 37°C for 18 to 24 hours. The diameter of the zones of inhibition were measured with a ruler and interpreted according to the guidelines provided by the National Committee for Clinical Laboratory Standards [16].

### Plasmid DNA extraction/Analysis

Plasmid extraction was performed by a simplified alkaline lyses method described by Birnboim and Doly (1979) [17]. Overnight culture (1.5 ml) was centrifuged at 5,000 rpm for 1 minute to pellet the cells. The supernatant was gently decanted. After washing, the cell pellet was re-suspended in 300 µl Tris (tris(hydroxymethyl)aminomethane) - EDTA (ethylenediaminetetraacetic acid) - NaOH-SDS (sodium dodecyl sulfate) (TENS) buffer solution and mixed by gentle inversion (five times) within 3–5 minutes of incubation on ice to obtain a straw-like, sticky lysate. This was followed by neutralization by adding 200 µl of 5 M potassium acetate buffer (pH 5.2), followed by incubation on ice for 10 minutes. Plasmid DNA solution was recovered as supernatant after centrifugation at 10,000 rpm for 10 minutes. Plasmid pellets isolated with ice-cold absolute ethanol were washed in 70% ethanol before re-suspension in 20–40 µl Tris (10 mM) - EDTA (1 mM) buffer (pH 8.0). The plasmid DNAs were separated by electrophoresis on 0.8% agarose pre-stained with ethidium bromide (0.5 µg/ml), the plasmid DNAs were visualized on high performance ultra violet transilluminator and photographed. Molecular weight of the plasmids were calculated based on the DNA maker.

### Plasmid eviction potential of plant extract

The plant eviction potential was determined by inoculating 18 hours old enteric pathogens harbouring plasmid DNA into a freshly prepared Trypton soya agar. The inoculum was spread evenly. The disc containing the plant was carefully placed in the middle of the medium and the plate was incubated at 37°C, for 18 hours. Zones of inhibition were observed. The organisms that fell between 2-3 mm from the periphery of the inhibition zone were scooped, and the process of plasmid isolation was repeated using the method described by Birnboim and Doly (1979) (17) for the presence or absence of plasmids. The loss of plasmid DNA was considered as plasmid eviction.

### Determination of minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC)

#### Broth dilution method

MIC and MBC of the effective extract was determined by broth dilution method. Equal volume of plant extract and basal medium (Tryptone soya broth) was mixed in series of test tubes. Standardized Inoculum 106CFU/ml was added to series of test tubes with the aid of a Micropipette. Incubation was carried out aerobically at 37°C for 18 to 24 hrs. Two control tubes were maintained for each test batch. These include tubes containing plant extract and growth medium without Inoculum (antibiotic control) and tube containing the medium, physiological saline and the Inoculum (organism control) in the test tubes. The lowest concentration (highest dilution) of the extract permitting no visible growth (any turbidity) when compared with the control tube was considered as the MIC. However, the MBC was determined by sub culturing the test dilution on a Tryptone soya agar and was further incubated for 18-24 hrs at 37°C. The lowest concentration that is highest dilution of the extract that yields no single bacterial colony on solid medium was regarded as MBC.

#### Phytochemical screening methods

**Test for alkaloids:** One ml of 1% Hydrochloric acid was added to 3ml of extract in a test tube. The mixture was heated for 20 minutes. It was cooled and filtered. Two drops of Mayer's reagent was added to the filtrate. [18]

**Test for tannins:** Half a gram of extract was added to 20 ml of distilled water. The suspension was properly mixed and filtered. 3 drops of 5% ferric chloride (FeCl<sub>3</sub>) was added to the filtrate. [18]

**Test for flavonoids:** One gram of extract was added to 10 ml of ethyl acetate and mixed. 1ml of diluted ammonia was added to the mixture [18].

**Test for saponins:** One gram of extract was added to 10 ml of distilled water. The mixture was shaken vigorously. [18]

**Anthraquinones:** Five gram of the extracts was added to 10 ml of ether and mixed. 1ml of ammonia solution was added to the filtrate. [18]

### Results

The profile of the three medicinal plants *Daniellia oliveri*, *Lannea welwitschii*, and *Lophira alata* used in this study is indicated in table 1. Antimicrobial activities of the crude plants extracts against the plasmid-carrying MDR bacterial strains are shown in table 2. At 5 to 40 mcg/ml concentration, the methanol extract of *L. Welwitschii*, recorded the highest zone of inhibition against all the strains of organisms used. The highest zone of inhibition recorded was against *S. Typhimurium* with a diameter of 14mm at 5mg/ml, 18.0mm at 20mcg/ml and 22mm at 40mcg/ml respectively. The zones of inhibition of the aqueous extracts against the organisms used were lower than the corresponding methanol extract. At 5mcg/ml, the crude extract of *Lophira alata* recorded no zone of inhibition against the strains of organisms used except for the standard strain *E.coli* ATCC25922 (Table 2). The MIC recorded in this study ranged from 0.625 to 20.0 and MBC ranged from 5.0 to 40.0. *Lannea welwitschii* recorded the lowest MIC and MBC values against the organisms used, while *Lophira alata* recorded the highest MIC and MBC values (Table 3).

The result of the phytochemical analysis of the plant extracts revealed the presence alkaloids, tannins, flavonoids in all the plant extracts used. Saponins were found in *D. oliveri* and *L. alata* while anthraquinone was found in both *D. oliveri* and *L. Welwitschii* (Table 4).

The results of the plasmid eviction potentials of the plants extracts are as shown in table 5. All the strains of *S. Typhimurium* had 100% loss rate. Significant plasmids loss was recorded in all the strains of organisms used except for two strains of *Klebsella* spp and a strain of *S. typhi* that had 0% loss rate (Figures 1 & 2).

**Table 1** Profile of the three medicinal plants used

Botanical name	Family name	Local name	Plant part used	Voucher number
<i>Daniellia oliveri</i> (Rolfe) Hutch & Dalz	Fabaceae	Ogbogbo	Stem	LSH1200
<i>Lannea welwitschii</i> (Hiern)	Anacardiaceae	Oriira	Stem	LSH1201
<i>Lophira alata</i> (Banks ex f. Gaertn.f)	Ochnaceae	Ponhan	Stem	LSH1202

**Table 2**Antibacterial activities of plants extracts on plasmid- carrying MDR enteric bacterial pathogens

Plant used	B a c t e r i a l p a t h o g e n s									
	<i>S. Typhi</i> (n-12)		<i>S. Typhimurium</i> (n-10)		<i>Shigella spp</i> (n-8)		<i>E. coli ATCC25922</i>		<i>Klebsiella spp</i> (n-10)	
	A E	M E	A E	M E	A E	M E	A E	M E	A E	M E
				5mcg/ml						
<i>D. oliveri</i>	-	-	-	6 . 0	6 . 0	8 . 0	6 . 0	8 . 0	-	-
<i>L. welwitschii</i>	8 . 0	10.0	10.0	14 . 0	-	6 . 0	8 . 0	12.0	8.0	10.0
<i>L. alata</i>	-	-	-	8 . 0	-	-	5 . 0	8 . 0	-	-

**20mcg/ml**

<i>D. oliveri</i>	6 . 0	8 . 0	-	8 . 0	8 . 0	12.0	12.0	14.0	-	-
<i>L. welwitschii</i>	12.0	16.0	14.0	18 . 0	-	8 . 0	14.0	18.0	10.0	14 . 0
<i>L. alata</i>	8 . 0	10.0	-	12 . 0	-	-	8 . 0	10.0	-	-

**40mcg/ml**

<i>D. oliveri</i>	10.0	12.0	8 . 0	14 . 0	12.0	14.0	14.0	18.0	-	10 . 0
<i>L. welwitschii</i>	14.0	20.0	18.0	22 . 0	8 . 0	12.0	18.0	24.0	12.0	18 . 0
<i>L. alata</i>	10.0	14.0	-	14 . 0	-	-	10.0	12.0	-	-
Ciprofloxacin10µl	16.0		14.0		10.0		18.0		12.0	

**Activity key:** figures indicate average diameter of zone of inhibition in mm

- = No inhibition, AE = aqueous extract, ME = methanol extract, N = number of isolate tested, Ciprofloxacin = commercial antibiotics used as control, *E. coli* ATCC 25922 = Standard control strain, MDR= multidrug resistant strain

**Table 3**Inhibitory and bactericidal concentration of plant extract on MDR- strains of enteric bacterial pathogens.

P l a n t	<i>S. Typhi</i>		<i>S. Typhimurium</i>		<i>Shigella spp</i>		<i>E. coli ATCC 25922</i>		<i>Klebsiella spp</i>	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
	I n m c g / m l									
<i>D. oliveri</i>	ND	ND	10.0	40.0	20.0	40.0	10.0	20.0	ND	ND
<i>L. welwitschii</i>	0.625	5.0	5.0	10.0	10.0	20.0	5.0	10.0	10.0	20.0
<i>L. alata</i>	20.0	40.0	20.0	ND	ND	ND	10.0	20.0	ND	ND

**Key:** MIC = minimum inhibitory concentration, MBC= minimum bactericidal concentration

ND = not determined, MDR = multidrug resistant strains

**Table 4** Plasmid analysis of the isolates before and after exposure to methanol extract of *Lannea welwitschii* at 10mcg/ml (MIC)

Lane/ SN	I S O L A T E	Number of plasmid before exposure	S i z e s , k b	Number of plasmid after exposure	Sizes ,kb	Lost rate
1	10 kb Plasmid DNA Maker	4	1 0 , 9 , 7 , 6			
2	<i>Salmonella</i> Typhi	1	25	1	25	0%
4	<i>Salmonella</i> Typhi	1	15	0	0	100%
6	<i>Salmonella</i> Typhi	4	25, 23, 6, 5	4	25, 23, 6, 5	0%
8	<i>Salmonella</i> Typhi	3	25, 23, 5	2	25, 23	33.3%
9	<i>Salmonella</i> Typhi	3	25, 23, 5	0	0	100%
14	<i>Salmonella</i> Typhi	4	10, 9, 7, 6	0	0	100%
17	<i>Salmonella</i> Typhi	1	5	0	0	100%
19	<i>Salmonella</i> Typhi	3	25, 23, 5	2	25, 23	33.3%
20	<i>Salmonella</i> Typhi	1	5	0	0	100%
21	<i>Salmonella</i> Typhi	1	5	0	0	100%
3	<i>Salmonella</i> Typhimurium	3	10, 9, 8	0	0	100%
4	<i>Salmonella</i> Typhimurium	2	10, 9	0	0	100%
5	<i>Salmonella</i> Typhimurium	2	9, 8	0	0	100%
7	<i>Salmonella</i> Typhimurium	3	9, 8, 7	0	0	100%
8	<i>Salmonella</i> Typhimurium	2	9, 8	0	0	100%
9	<i>Salmonella</i> Typhimurium	3	9, 8, 5	0	0	100%
10	<i>Salmonella</i> Typhimurium	3	10, 9, 8	0	0	100%
12	<i>Salmonella</i> Typhimurium	3	9, 8, 7	0	0	100%
1	<i>Shigella</i> sp	2	23, 15	1	23	50%
2	<i>Shigella</i> sp	1	10	0	0	100%
3	<i>Shigella</i> sp	1	9	0	0	100%
5	<i>Shigella</i> sp	2	12, 15	0	0	100%
7	<i>Shigella</i> sp	3	15, 10, 8	1	15	66.7%
8	<i>Shigella</i> sp	1	10	0	0	100%
9	<i>Shigella</i> sp	1	7	0	0	100%
11	<i>Shigella</i> sp	2	12, 10	0	0	100%
12	<i>Klebsiella</i> sp	1	8	0	0	100%
13	<i>Klebsiella</i> sp	1	15	0	0	100%
14	<i>Klebsiella</i> sp	3	25, 20, 10	2	25, 20	33.3%
16	<i>Klebsiella</i> sp	2	15, 8	0	0	100%
17	<i>Klebsiella</i> sp	1	14	0	0	100%
19	<i>Klebsiella</i> sp	1	12	0	0	100%
21	<i>Klebsiella</i> sp	1	12	0	0	100%
24	<i>Klebsiella</i> sp	2	10, 8	0	0	100%
25	<i>Klebsiella</i> sp	1	25	1	25	0%
26	<i>Klebsiella</i> sp	2	23, 15	2	23, 15	0%

**Table 5** Phytochemical analysis of the three medicinal plant extracts used

P l a n t   s c r e e n e d		C o m p o n e n t s									
Plant species	Plant part	Alkaloids		T a n n i n s		Flavonoids		Saponins		Anthraquinones	
		ME	AE	ME	AE	ME	AE	ME	AE	ME	AE
<i>D. oliveri</i>	S t e m	+	+	+	+	+	+	+	+	+	+
<i>L. welwitschii</i>	S t e m	+	+	+	+	+	+	-	-	+	+
<i>L. alata</i>	S t e m	+	+	+	+	+	+	+	+		

**Key:** +++ Appreciable amount

++ Moderate amount

+ Trace amount


- Complete absence

ME: Methanol extract


AE: Aqueous extract



**Figure 1:** Representation of Plasmid analysis before the introduction of the extract of *Lannea welwitschii*. Lane 1=10 kb Plasmid DNA Maker, lane 2=25kb *S. Typhi*, lane 3= *S. Typhi*, lane 4=15kb *S. Typhi*, lane 5= *S. Typhi*, lane 6=25, 23.6, 5 kb *S. Typhi*, lane 7= *S. Typhi*, lane 8=25, 23, 5kb *S. Typhi*, lane 9= 25, 23, 5kb lane 10= *S. Typhi*, lane 11-13= *S. Typhi*, lane 14=10, 9, 7 *S. Typhi*, lane 15-16= *S. Typhi*, lane 17= 5kb lane 18= *S. Typhi*, lane 19= 23,4kb *S. Typhi*, lane 20= 5kb *S. Typhi*, lane 21= 5kb *S. Typhi*.



**Figure 2:** Representation of plasmid Analysis after the introduction of the extract of *Lannea welwitschii*. Lane 1=10 kb Plasmid DNA Maker, lane 2=*S. Typhi*, lane 3= *S. Typhi*, lane 4= *S. Typhi*, lane 5=Shigella spp, lane 6= Shigella spp, lane 7= Shigella spp, lane 8=25, 23kb *S. Typhi*, lane 9= 25kb *Klebsiella* spp, lane 10= *Klebsiella* spp, lane 11= 25kb *S. Typhi*, lane 12= *Klebsiella* spp, lane 13= *Klebsiella* spp, lane 14= 10, 9, 7. 5kb *S. Typhi*, lane 15= Shigella spp, lane 16= Shigella spp, lane 17=25, 23, 6, 5kb *S. Typhi*, lane 18= 5kb *S. Typhi*, lane 19= 5kb *S. Typhi*, lane 20=Shigella spp, lane 21=Klebsiella spp, lane 22=Klebsiella spp, lane 23=Klebsiella spp, lane 24= 25, 23kb *S. Typhi*, lane 25= 25kb *Klebsiella* spp, lane 26= *Klebsiella* spp.



## Discussion

Medicinal plants are considered as a potential source of new chemotherapeutic drugs because of their diverse phytochemicals. In this study antimicrobial activities of the crude plant extracts against the plasmid-carrying MDR bacterial strains were carried out at 5 to 40 mcg/ml concentration. The methanolic extract of *L.*

*welwitschii* at 10mg/ml was found to be active against strains of *S. Typhi*, *S. Typhimurium*, *E. coli*, *Shigella* spp and *Klebsiella* spp. High zone of inhibition was recorded against *S. Typhimurium* with a diameter of 14mm at 5mcg/ml, 18.0mm at 20mcg/ml and 22mm at 40mcg/ml respectively. This result was consistent with previous studies conducted elsewhere where increase in the antimicrobial activities of the plant extract against the MDR organisms were reported to be attributable to the increase in the concentration and the nature of the active components. Also, that the active components are water soluble and are easily extracted by water [15,19,20]. Traditionally, the plant is soaked in water for days, thereafter ready for drinking for the treatment of patients with gastro-intestinal disorder. Our findings therefore further corroborated the preference of water extract of the plants by traditional medical practitioners. Generally, the MIC of the three medicinal plants evaluated in this study ranged from 0.625 to 20.0 mcg/ml and MBC from 5.0 to 40.0mcg/ml. The MIC values of the three extracts were lower than the MBC value indicating that, these plants are bacteriostatic at lower concentration and bactericidal at higher concentration, an assertion that had been documented by several workers [9,15,21]. Inactivity of some of the extracts at lower concentration was observed in this study. This was probably attributed to the presence of plasmid in most of the organisms used. In this study, no zone of inhibition was recorded against MDR-enteric pathogens tested at 5mcg/ml concentration of the crude extract of *Lophira alata*. This however, indicated complete inactivity of the extract except for the standard strain of *E. coli* ATCC25922 used, where certain level of activity was recorded. Similarly, both the aqueous and methanol extract of *D. oliveri* and *L. alata* exerted little activity against most of the strains of MDR enteric pathogens tested. It should be noted that *L. alata* is traditionally used for the treatment of typhoid fever and *D. oliveri* for the treatment of dysentery in Abeokuta, Ogun state Nigeria. The implication of this is that, these two medicinal plants are not effective for the treatment of MDR-enteric bacterial-associated diseases, thus their continuous usage should be discouraged.

The phytochemical analysis of the aqueous and methanol extracts of *L. welwitschii*, *D. Oliveri* and *L. alata* showed the presence of bioactive compounds such as alkaloid, saponins, tannins, flavonoids and anthraquinones which had been documented to inhibit intestinal mobility in a dose related manner [12,19,22]. The extract of *L. welwitschii* has also been reported by Amole et al., [12] to produce a movement at standard charcoal meal in the small intestine, suggesting an antispasmodic activity. The effect of *L. welwitschii* against enteric pathogens as observed in this study may be due to the presence of appreciable amount of tannins, anthraquinones and flavonoids as indicated in the result of the phytochemistry. Flavonoids had been reported to be hydroxylated phenolic substances linked to an aromatic ring and are known to be synthesized by plants in response to microbial infection [23]. Their activity is probably due to their ability to complex with extracellular and soluble proteins and to complex with bacterial cell walls. Flavonoids, also have been reported to disrupt microbial membranes [24].

The plasmid eviction potentials of the plant extracts were evaluated. All the strains of *S. Typhimurium* had 100% loss rate. For example *Shigella* spp had 100% loss rate. Seven out of the ten strains of *Klebsella* spp had 100% loss rate. Generally, significant plasmids loss was recorded in all the strains of organisms used except for two strains of *Klebsella* spp and *S. Typhi* with no antiplasmidic effect observed because of 0% loss rate of Plasmid DNA. (Figures 1 & 2) It was also observed that high molecular weight plasmid DNA was not evicted (Table 4). Nevertheless, antimicrobial activity and DNA intercalation properties of the active constituents had been documented in literature [25]. The curing ability of this plant extract might be due to its active constituent, which may intercalate with the DNA molecule and might have inhibited plasmid replication selectively at sub-MIC concentration (below the MIC of 10 mcg/ml used). Curing ability of some medicinal plants on plasmid-encoded MDR- pathogens had been reported [26-28]. Also, plant derived curing agents; Plumbagin, 5-hydroxy-2-methyl-1, 4-naphthoquinone isolated from *Plumbago zeylanica* had been reported to cure R-plasmids in *E. coli* [29]. In another study by Beg et al (2004) [30], the alcoholic extract of *P. zeylanica* cured R-plasmid harbouring *E. coli* with 14 per cent. Recently antiplasmid activity of essential oils was reported by Schelz et al 92006 [31]. The antiplasmidic activity of medicinal plants seems to be a possible novel effective curing agents that may pave the way for further drug research. Therefore the antiplasmidic effect of *L. welwitschii* observed in this study calls for further investigation, as this plant may be a useful natural resource, for the development of novel drug that will complement the orthodox antimicrobials against increasing emerging trends of MDR bacterial pathogens.

## Conclusion

This study revealed that methanol extracts of the three medicinal plants were active against enteric bacterial pathogens. It was observed that extracts of *L. alata* and *D. oliveri* were not active against MDR-enteric bacterial-associated diseases, thus their continuous usage should be discouraged. Also crude methanolic extract of *L. welwitschii* exerted antiplasmidic effect on multiple drug resistant enteric bacterial pathogens. The present result has offered organic extracts of *L. welwitschii* as a plasmid curing agent. This finding suggests possibility of a new approach in drug formulation using anti-plasmidic plant extracts for effective management of plasmid encoded drug resistant enteric bacteria-associated diseases. Further studies are required using modern methods for characterizing resistance elements to assess the effects of extract of *L. welwitschii* on clinically significant enteric bacteria.

## Conflict of interest statement

The Author declared that there is no conflict of interest

## ACKNOWLEDGEMENTS

## References :

1. WHO: World health organization guideline on good agricultural and collection practices (GACP) of medical plants 2003, P80
2. Menashe O, Kagansaya E, Baasov T, Yaron S: Aminoglycoside affect intracellular *Salmonella enterica* serovar Typhimurium and Virchow. *Antimicrobial Agent Chemother*, 2008, 52(3):920-6 doi 1128/AAC.00382-02
3. Faleyimu OI, Akinyemi O, Idris YM: Survey of forest plant used in traditional treatment of typhoid fever in Chikun local government area of Kaduna state, Nigeria: *International journal biomedical science*. 2010, 6:75-81
4. Fabeku PO: Traditional medicine: the art, ways and practices in: Odugbemi T editor, *Outline and pictures of medicinal plants from Nigerian*. University of Lagos press; 2006, P13-24
5. Newman RR, it Hrich E: Commercialization of non-timber forest products: review and analysis for research, Indonesia: CIFOR 2000, P176
6. Adaku VIW, Okewesili FCN: Anti hyperglycaemic effect of aqueous extract of *Daniellia oliveri* and *Sarcocephalus latifolius* roots on key carbohydrate metabolic enzymes and glycogen in experimental diabetes. *BIOKEMISTRI* 2008, 20(2):63-70
7. Ahmadu, A., Haruna, A.K., Magaji, G., Yaro, A.H (2008). Antidiarrheal activity of the leaves of *Daniellia oliveri* Hutch and Daiz (Fabaceae) and *Ficus syncomorus* MIQ (Moraceae), *African journal of traditional, complementary and alternative medicines* 2007, 4:524-528
8. Ainstie JA: List of plants used in native medicine in Nigeria. Oxford university imperial forest institute paper T. 1990. Cited Irvine 1961.
9. Yahaya O, Tijani MB, Abraham OJ, Umar JO, Maichi OE, Usman B: Antibacterial activity of the crude extract of *Daniellia oliveri* against some bacteria associated with enteric infections, *J of medical Applied Bioscience* 2011, 3:26-31
10. Adegoke EA, Akinsanya A, Nagri SHZ: Studies of Nigerians medicine plant: A preliminary survey of plant alkaloids. *J. West Africa Sci, Assos* 1968, 73:13-17
11. Iwu, MM, Unaeze NC, Okunji CO, Corley DG, Samson DR, Tempesta MS (1991). Antimicrobial activity of isothiocyanates from the essential oil of *Hippocrates welwitschii* roots. *J of Pharmaceutical Biology*, 1991, 29[2]:154-158 Doi: 10.3109/13880209109082869
12. Amole OO, Salahdeen HM, Onyiahialam A: Evaluation of antidiarrheal effect of *Lannea welwitschii* hiern (Anacardiaceae) bark extract. *African journal of pharmacy and pharmacology* 2010, 4(4) 165-169
13. McDonald I, Joseph O, Harriet M: Documentation on medicinal plant sold in market in Abeokuta, Nigeria. *Tropical journal of pharmaceutical research*. 2010, 9(2):110-118
14. Odugbeni T. Medicinal plant species, family names and uses: A text book of medicinal plant. Published by the university press of Lagos, Nigeria, 2008
15. Akimiyemi KO, Mendie UE, Smith ST, Oyefolu, AO, Coker, AO: Screening of some medicinal plants used in south west Nigeria traditional medicine for Anti-salmonella typhi activity. *J of herbal Pharmacotherapy* 2005, 5 (1):45-60
16. National committee for clinical laboratory standards: Performance standard for antimicrobial susceptibility testing; 2002, twelfth informational supplement, Wayne, PA, USA

17. Birnboim HC, Doly J: A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acid Res* 1979, 7:1513-1524.
18. Trease GE, Evans WC: *Pharmacognosy*. 15th edition. Harcourt publisher; Edinburgh. 2002, Pp334-337
19. Adebayo GB, Oguntoye SO, Agbowo EU: Phytochemical analysis and the antibacterial screening of the plant *Daniellia oliveri* stem bark. *Chem. Dept. University of Ilorin: CSN Conference proceeding* 2010:1-9
20. Dickson AM, Fred OCN, Ganiyu OY: A comparative study of aqueous, methanol and chloroform extracts of some selected medicinal plants used in Igala land of Nigeria. *Der pharmacia sinica*. 2011, 2(1):222-227
21. Sule LO, Agbabiaka TO, Oyeyiola GP: Traditional values of cotton leaf and lemon juice against clinical bacterial isolates. *Ethnobotanical leaflets* 2009, 13:522-31
22. Carlo DC, Mascolo N, Izzo AA, Capasso F: Effect of quercetin on the gastro intestinal tract in rats and mice. *Phytol Therapy Res* 1994, 6:42-45
23. Marjorie MC: Plant products as antimicrobial agents. *Clinical microbiology review* 1999, 564-582
24. Tnchiya H, Sato M, Miyazaki T, Tanigaki S, Obyama M, Tanaka T, Iinuma M: Comparative study on the antimicrobial activities of phytochemicals flavanones against methicillin-resistant *Staphylococcus aureus* J. *Ethnopharmacol*. 1996, 50:27-34
25. Bailly C, Laine W, Baldeyrou B, et al: DNA intercalation, topoisomerase II inhibition and cytotoxic activity of plant alkaloid neocryptolepine. *Anticancer drug Des*. 2000, 15(3):191-201

## References :

26. Adel K, Srwa M: Potential of aqueous and alcohol extracts of *Quercus infectoria*, *Linum usitatissimum* and *Cinnamomum zeylanicum* as antimicrobials and curing of antibiotic resistance in *E. coli*. *Current Research Journal of Biological Sciences* 2010, 2(5): 333-337
27. Jasmine RS, Selvakumar, BN: Synergistic action of two Indian medicinal plants on clinical isolates of vancomycin resistant enterococci species. *Pharmacologyonline, Newsletter*, 2011, 2: 898-904
28. Varsha S, Vinay K, Javed M, Latha. C: Curing of Plasmid-Mediated Antibiotic Resistance in Multi-Drug Resistant Pathogens Using *Alpinia galanga* Rhizome Extract. *Adv Bio Tech*: 2013, 13(01): 01-05.
29. Lakshmi VV, Padma S, Polasa H: Elimination of multidrug-18. resistant plasmid in bacteria by plumbagin, a compound derived from a plant. *Curr Microbiol* 1987, 16: 159-61.
30. Beg AZ, Ahmad I: Effect of *Plumbago Zeylinica* extract and certain curing agents on multidrug resistance bacteria of clinical origin. *World J Microb Biotech* 2004, 16: 841-4.